



Contribution ID: 28

Type: Review Talk

Analytical laboratory studies of solar system dust

Wednesday 13 June 2018 14:45 (45 minutes)

Contemporary solar system dust is collected as interplanetary dust particles (IDPs) and micrometeorites (MMs) in the stratosphere using aircraft, and at polar and mid-latitude ground locations using several sampling methods. The collections are fundamentally important to cosmochemistry, planetary science and early solar system accretional processes in particular because some IDPs and MMs are from parent bodies, including comets and other small, icy bodies, that are not well represented among known meteorite groups [1]. Small bodies are more likely to contain well-preserved solids from the outer solar nebula and presolar environments. The minuscule masses of individual IDPs and MM have severely limited analytical measurements but advances in instrumentation are expanding the scope of measurements and providing fundamental insight. It is well established that IDPs and MM contain the highest abundances of presolar refractory constituents (e.g. crystalline oxides and silicates) and recent studies have identified a population of non-refractory “soft” constituents [2]. All IDPs and MM are pulse heated, many to incandescence, for several seconds during atmospheric entry followed by exposure to terrestrial contamination. A key finding and challenge arising from the study of “soft” constituents are that well-preserved IDPs and MM appear to be much rarer than has previously been assumed.

REFERENCES:

1. S. Taylor et al. (2016). Elements, 12(3).
2. H. A. Ishii, et al (2018), Proc. Natl. Acad. Sci., in press.

Consider for a poster?

Author: Dr BRADLEY, John (University of Hawaii)

Presenter: Dr BRADLEY, John (University of Hawaii)

Session Classification: Dust in the solar system

Track Classification: Dust in planet-forming disks