Cosmic Dust: origin, applications & implications



Contribution ID: 36 Type: Poster

Temperature-Dependent Laboratory Measurements of the Far-Infrared to Millimeter Opacity of Carbonaceous Dust-Analogues

Monday, 11 June 2018 10:48 (1 minute)

We are measuring and analysing the FIR- and THz- Spectra of pyrolysed micro-crystalline cellulose as an analogue of carbonaceous interstellar dust. We are using cellulose-powder with crystal sizes of about 20\, μ m and are heating it up to 1000° C. First results of the mass normalised extinction are presented and compared to J\"ager et al. (1998). The temperature dependent measurements took place in a dry environment at room temperature (RT) down to the environmental temperature of $T_{\rm env} = 10$ \,K. \\

Our aim is to assess carbonaceous dust analogues in terms of structure, nature and morphology. For theoretical and observational investigation we are going to determine their optical constants. Furthermore, we are going to calculate the emission cross section of particles with different geometries to compare them with the measured results.

Consider for a poster?

Yes

Primary author: GREIF, Jonas (Astrophysical Institute and University-Observatory Jena)

Co-author: Dr MUTSCHKE, Harald (Astronomical Institute and University-Observatory Jena)

Presenter: GREIF, Jonas (Astrophysical Institute and University-Observatory Jena)

Session Classification: Poster Presentations

Track Classification: What is dust?