



Contribution ID: 35

Type: **Contributed talk**

Spherical harmonics expansion: excellent tool for analysis of anisotropic small-angle scattering data

With small-angle scattering, we may study the structure of (soft) matter, e.g. polymer melts, on length scales of ~1-100 nm. If there is alignment in the system on these length scales, the small-angle scattering pattern will be anisotropic.

The anisotropy of the scattering patterns can be quantified by expansion in spherical harmonics which separates contributions based on their symmetries.

We demonstrate the power of the spherical harmonics expansion framework [1], by analyzing small-angle neutron scattering data for the relaxation of a polymer melt following fast uniaxial extension, previously published in [2]. We see that the chains initially retract which was proposed in 1978 [3] and has been debated since [1, 4-7].

References

- [1] Z. Wang et al, Phys Rev X 7, 031003 (2017)
- [2] J.J.K. Kirkensgaard et al, Phys. Rev. E 94, 020502(R) (2016)
- [3] Doi and Edwards, J. Chem. Soc., Faraday Trans. 2, 74, 1802-1817 (1978)
- [4] Boue et al, Journal de Physique, 43, 137-148 (1982)
- [5] Mortensen et al, MRS Proceedings, 79, 259 (1986)
- [6] Blanchard et al, Phys. Rev. Lett., 95, 166001 (2005)
- [7] Hsu and Kremer, Phys. Rev. Lett., 121, 167801 (2018)

Author: Ms BORGER, Anine (University of Copenhagen, Niels Bohr Institute)

Co-authors: KIRKENSGAARD, Jacob (Niels Bohr Institute, University of Copenhagen); MORTENSEN, Kell (University of Copenhagen); Prof. ALMDAL, Kristoffer (Technical University of Denmark, Department of Micro- and Nanotechnology); Prof. HASSAGER, Ole (Technical University of Denmark, Department of Chemical and Biochemical Engineering); Dr HUANG, Qian (Technical University of Denmark, Department of Chemical and Biochemical Engineering)

Presenter: Ms BORGER, Anine (University of Copenhagen, Niels Bohr Institute)