## Coaction for Feynman Integrals

#### Ruth Britto

Trinity College Dublin

Amplitudes 2021 August 20, 2021

Based on 2106.01280 with Samuel Abreu, Claude Duhr, Einan Gardi, and James Matthew







Trinity College Dublin Coláiste na Tríonóide, Baile Átha Cliath The University of Dublin

<ロ> (日) (日) (日) (日) (日)

3

We conjecture three compatible coactions on Feynman integrals:

• Local coaction on multiple polylogarithms (MPLs), elliptic multiple polylogarithms, etc.

This one is well known. [Goncharov, Brown]

Applies in the Laurent expansion of Feynman integrals in the parameter  $\epsilon$  of dimensional regularization.

• Global coaction on generalized hypergeometric functions.

Applies to Feynman integrals in dimensional regularization without taking the Laurent expansion.

Conjectured to exist for arbitrary Feynman integrals; examples found with integer-based parameters; [Abreu, RB, Duhr, Gardi, Matthew] since proven for Lauricella functions [Brown, Dupont].

#### • Diagrammatic coaction

1st claim: the output from the other two coactions are compatible with each other and can be repackaged as Feynman integrals.

2nd claim: the coaction output can also be obtained by applying graphical operations before evaluating.

- Coaction is naturally compatible with discontinuities and differential operators, so we hope it can be applied to new computations
- Dimensional regularization is essential
- Formally, we distinguish motivic and de Rham MPLs, hypergeometric functions, etc. or use single-valued versions
- This talk: general principles and 1- and 2-loop examples

$$\underbrace{\frac{1}{e_1}}_{e_1} \underbrace{e_2}_{e_1} = \frac{e^{\gamma_E \epsilon} \Gamma(1+\epsilon)}{\epsilon(1-\epsilon)} (m^2)^{-1-\epsilon} {}_2F_1\left(1, 1+\epsilon; 2-\epsilon; \frac{p^2}{m^2}\right)$$

$$= \frac{1}{p^2} \left[ \frac{\log\left(\frac{m^2}{m^2-p^2}\right)}{\epsilon} + \operatorname{Li}_2\left(\frac{p^2}{m^2}\right) + \log^2\left(1-\frac{p^2}{m^2}\right) + \log(m^2)\log\left(1-\frac{p^2}{m^2}\right) \right] + \mathcal{O}(\epsilon)$$

$$\frac{1}{e_1} \left( e_2 - \frac{e^{\gamma_E \epsilon} \Gamma(1+\epsilon)}{\epsilon_1} (m^2)^{-1-\epsilon} {}_2F_1\left(1, 1+\epsilon; 2-\epsilon; \frac{p^2}{m^2}\right) \right)$$

$$= \frac{1}{p^2} \left[ \frac{\log\left(\frac{m^2}{m^2-p^2}\right)}{\epsilon} + \operatorname{Li}_2\left(\frac{p^2}{m^2}\right) + \log^2\left(1-\frac{p^2}{m^2}\right) + \log(m^2)\log\left(1-\frac{p^2}{m^2}\right) \right] + \mathcal{O}(\epsilon)$$

$$\begin{array}{lll} \Delta(\log z) &=& 1 \otimes \log z + \log z \otimes 1 \\ \Delta(\log^2 z) &=& 1 \otimes \log^2 z + 2 \log z \otimes \log z + \log^2 z \otimes 1 \\ \Delta(\operatorname{Li}_2(z)) &=& 1 \otimes \operatorname{Li}_2(z) + \operatorname{Li}_2(z) \otimes 1 + \operatorname{Li}_1(z) \otimes \log z \end{array}$$

Ruth Britto, Trinity College Dublin

$$= \frac{1}{e_1} \left[ e_3 = \frac{e^{\gamma_E \epsilon} \Gamma(1+\epsilon)}{\epsilon(1-\epsilon)} (m^2)^{-1-\epsilon} {}_2F_1\left(1, 1+\epsilon; 2-\epsilon; \frac{p^2}{m^2}\right) \right]$$

$$= \frac{1}{p^2} \left[ \frac{\log\left(\frac{m^2}{m^2-p^2}\right)}{\epsilon} + \operatorname{Li}_2\left(\frac{p^2}{m^2}\right) + \log^2\left(1-\frac{p^2}{m^2}\right) + \log(m^2)\log\left(1-\frac{p^2}{m^2}\right) \right] + \mathcal{O}(\epsilon)$$

$$\begin{split} \Delta\Big({}_{2}F_{1}(\alpha,\beta;\gamma;x)\Big) &= {}_{2}F_{1}(1+a\epsilon,b\epsilon;1+c\epsilon;x) \otimes {}_{2}F_{1}(\alpha,\beta;\gamma;x) \\ &- \frac{b\epsilon}{1+c\epsilon} {}_{2}F_{1}(1+a\epsilon,1+b\epsilon;2+c\epsilon;x) \\ &\otimes \frac{\Gamma(1-\beta)\Gamma(\gamma)}{\Gamma(1-\beta+\alpha)\Gamma(\gamma-\alpha)} x^{1-\alpha} {}_{2}F_{1}\left(\alpha,1+\alpha-\gamma;1-\beta+\alpha;\frac{1}{x}\right) \end{split}$$

Coaction for Feynman Integrals

$$= \frac{1}{e_1} \left[ e_3 = \frac{e^{\gamma \varepsilon \epsilon} \Gamma(1+\epsilon)}{\epsilon(1-\epsilon)} (m^2)^{-1-\epsilon} {}_2F_1\left(1,1+\epsilon;2-\epsilon;\frac{p^2}{m^2}\right) \right]$$
$$= \frac{1}{p^2} \left[ \frac{\log\left(\frac{m^2}{m^2-p^2}\right)}{\epsilon} + \operatorname{Li}_2\left(\frac{p^2}{m^2}\right) + \log^2\left(1-\frac{p^2}{m^2}\right) + \log(m^2)\log\left(1-\frac{p^2}{m^2}\right) \right] + \mathcal{O}(\epsilon)$$



Coaction for Feynman Integrals

イロト イヨト イヨト イヨト

æ

## Preview of 2-loop example

Sunset with two massive propagators.



General formula for coaction on integrals:

$$\Delta\left(\int_{\gamma}\omega\right)=\sum_{i}\int_{\gamma}\omega_{i}\otimes\int_{\gamma_{i}}\omega$$

Ruth Britto, Trinity College Dublin

Coaction for Feynman Integrals

イロン 不同 とくほど 不同 とう

э

## General formula for coaction on integrals

$$\Delta\left(\int_{\gamma}\omega
ight)=\sum_{i,j}\mathsf{c}_{ij}\int_{\gamma}\omega_{i}\otimes\int_{\gamma_{j}}\omega$$

- All three coactions have this structure.
- Satisfies axioms of coaction.
- Claim of this formula: there exist sets  $\{\omega_i\}$ ,  $\{\gamma_j\}$ ,  $\{c_{ij}\}$  to make it true.
- {ω<sub>i</sub>} generate cohomology
- $\{\gamma_j\}$  generate homology
- {c<sub>ij</sub>} are rational in ε, algebraic in other parameters/kinematic variables; uniquely fixed by choices of {ω<sub>i</sub>} and {γ<sub>j</sub>}

(日) (部) (注) (注) (三)

## General formula for coaction on integrals

$$\Delta\left(\int_{\gamma}\omega\right)=\sum_{i,j}c_{ij}\int_{\gamma}\omega_{i}\otimes\int_{\gamma_{j}}\omega$$

Choice of bases:

- $\{\omega_i\}$ Left entries  $\int_{\gamma} \omega_i$  related to  $\int_{\gamma} \omega$  by standard IBP reduction. Choose them to be pure.
- $\{\gamma_j\}$

Right entries  $\int_{\gamma_j} \omega$  related to  $\int_{\gamma} \omega$  by change of contour. Can start with all possible cuts, but there are relations among them.

An important relation for 1-loop integrals:

$$\sum_{i} C_i I_n + \sum_{i < j} C_{ij} I_n = -\epsilon I_n \mod i\pi$$

applies to subgraphs of multiloop graphs.

Principle: no uncut loops needed in right entries. Use only "genuine L-loop cuts."

Ruth Britto, Trinity College Dublin

Coaction for Feynman Integrals

### Dual bases

Compact version of master formula:

$$\Delta\left(\int_{\gamma}\omega\right)=\sum_{i}\int_{\gamma}\omega_{i}\otimes\int_{\gamma_{i}}\omega$$

Look for bases such that

$$\int_{\gamma_j} \omega_i = \delta_{ij} + \mathcal{O}(\epsilon)$$

## Principles of the diagrammatic coaction

$$\Delta\left(\int_{\gamma}\omega
ight)=\sum_{i,j}{m{c}_{ij}\int_{\gamma}\omega_i\otimes\int_{\gamma_j}\omega}$$

- Left entries related by IBP, only L-loop diagrams
- Right entries are genuine *L*-loop cuts
- Coefficients  $c_{ij}$  are explicit for L = 1, [Abreu, RB, Duhr, Gardi] ad hoc for L > 1 as of now.
- Can dualize bases either before or after initial choice
- Consistent with degenerate limits
- UV/IR pole cancellation
- $\gamma$  can be a cut contour. Gives coaction on cut integrals.

・ロト ・四ト ・ヨト ・ヨト

### First 2-loop example: 1-mass sunset



$$S(\nu_1, \nu_2, \nu_3, \nu_4, \nu_5; D; p^2, m^2) = \left(\frac{e^{\gamma_E \epsilon}}{i\pi^{D/2}}\right)^2 \int d^D k \, d^D l \frac{[(k+l)^2]^{-\nu_4}[(l+p)^2]^{-\nu_5}}{[k^2]^{\nu_1}[l^2]^{\nu_2}[(k+l+p)^2 - m^2]^{\nu_3}}$$

2 master integrals, normalized to  $1 + \mathcal{O}(\epsilon)$ .

$$\begin{split} S^{(1)} &= \epsilon^2 \left( p^2 - m^2 \right) S(1, 1, 1, 0, 0; 2 - 2\epsilon; p^2, m^2) \\ &= (m^2)^{-2\epsilon} \left( 1 - \frac{p^2}{m^2} \right) e^{2\gamma_E \epsilon} \Gamma(1 + 2\epsilon) \Gamma(1 - \epsilon) \Gamma(1 + \epsilon) {}_2F_1 \left( 1 + 2\epsilon, 1 + \epsilon; 1 - \epsilon; \frac{p^2}{m^2} \right) \\ S^{(2)} &= -\epsilon^2 S(1, 1, 1, -1, 0; 2 - 2\epsilon; p^2, m^2) \\ &= (m^2)^{-2\epsilon} e^{2\gamma_E \epsilon} \Gamma(1 + 2\epsilon) \Gamma(1 - \epsilon) \Gamma(1 + \epsilon) {}_2F_1 \left( 2\epsilon, \epsilon; 1 - \epsilon; \frac{p^2}{m^2} \right) \end{split}$$

Hence only two 2 independent integration contours, e.g. the maximal cuts.

Ruth Britto, Trinity College Dublin

Coaction for Feynman Integrals

・ロト ・ 日 ト ・ ヨ ト ・ ヨ ト ・ ヨ

#### First 2-loop example: 1-mass sunset

Maximal cut integral:

$$C_{123}S^{(1)} \sim \int dk_0 \ k_0^{-1-2\epsilon} \left(p^2 - m^2 + 2\sqrt{p^2}k_0\right)^{-1-2\epsilon} \left(p^2 + 2\sqrt{p^2}k_0\right)^{2\epsilon}$$

After taking three residues, there is one integration left, of the hypergeometric form  $_2F_1$ .

Can choose two independent contours:

$$\Gamma_{123}^{(1)}: k_0 \in \left[-\frac{\sqrt{p^2}}{2}, 0\right] \qquad \Gamma_{123}^{(2)}: k_0 \in \left[\frac{m^2 - p^2}{2\sqrt{p^2}}, 0\right]$$

Results:

$$\begin{split} &\int_{\Gamma_{123}^{(1)}} \omega^{(1)} = 2\epsilon \ e^{2\gamma_E \epsilon} \frac{\Gamma(1+\epsilon)\Gamma(1-\epsilon)}{\Gamma(1-2\epsilon)} (p^2 - m^2)^{-2\epsilon} {}_2F_1\left(-2\epsilon, 1+2\epsilon; 1-\epsilon; \frac{p^2}{p^2 - m^2}\right) \\ &\int_{\Gamma_{123}^{(2)}} \omega^{(1)} = 4\epsilon \ e^{2\gamma_E \epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-4\epsilon)} (p^2)^{2\epsilon} (p^2 - m^2)^{-4\epsilon} {}_2F_1\left(-2\epsilon, -\epsilon; -4\epsilon; 1-\frac{m^2}{p^2}\right) \end{split}$$

Ruth Britto, Trinity College Dublin

Coaction for Feynman Integrals

## First 2-loop example: 1-mass sunset

Can arrive at compact/dual form  $\Delta \left( \int_{\gamma} \omega \right) = \sum_{i} \int_{\gamma} \omega_{i} \otimes \int_{\gamma_{i}} \omega$  by requiring  $\int_{\gamma_{j}} \omega_{i} = \delta_{ij} + \mathcal{O}(\epsilon)$ . Solution:

$$\gamma_{123}^{(1)} = \frac{1}{4\epsilon} \Gamma_{123}^{(2)}, \qquad \gamma_{123}^{(2)} = \frac{1}{2\epsilon} \left( \Gamma_{123}^{(1)} - \frac{1}{2} \Gamma_{123}^{(2)} \right).$$

$$\begin{split} \Delta S^{(1)} &= S^{(1)} \otimes \mathcal{C}_{123}^{(1)} S^{(1)} + S^{(2)} \otimes \mathcal{C}_{123}^{(2)} S^{(1)} \\ \Delta S^{(2)} &= S^{(1)} \otimes \mathcal{C}_{123}^{(1)} S^{(2)} + S^{(2)} \otimes \mathcal{C}_{123}^{(2)} S^{(2)} \end{split}$$





Check agreement with  $\Delta[_2F_1]$ .

Ruth Britto, Trinity College Dublin

・ロン ・回 と ・ ヨン ・ ヨン

#### 1-mass sunset: comments on cuts

•  $\gamma_{123}^{(1)}$  and  $\gamma_{123}^{(2)}$  generate full homology, including uncut contour

$$\int_{\Gamma_0} \omega^{(i)} = \mathbf{a} \int_{\gamma_{123}^{(1)}} \omega^{(i)} + \mathbf{b} \int_{\gamma_{123}^{(2)}} \omega^{(i)} \mod i\pi$$



Discontinuities can be recovered

$$\begin{aligned} \operatorname{Disc}_{m^2} S^{(i)} &\sim 2\epsilon \left( \mathcal{C}_{123}^{(1)} S^{(i)} - \mathcal{C}_{123}^{(2)} S^{(i)} \right) \\ \operatorname{Disc}_{p^2} S^{(i)} &\sim -4\epsilon \, \mathcal{C}_{123}^{(1)} S^{(i)} \end{aligned}$$

• Coaction takes the same form on cut integrals



## Diagrammatic coaction at two loops



 $\Delta(fg) = (\Delta f) (\Delta g)$ 

Ruth Britto, Trinity College Dublin

Coaction for Feynman Integrals

<ロ> <回> <回> < 回> < 回> < 回> < 回> = 三

$$\begin{split} S^{(1)} &= -\epsilon^2 e^{2\gamma_E \epsilon} \int \frac{d^{2-2\epsilon} k}{i\pi^{1-\epsilon}} \int \frac{d^{2-2\epsilon} l}{i\pi^{1-\epsilon}} \frac{\sqrt{\lambda \left(p^2, m_1^2, m_2^2\right)}}{(k^2 - m_1^2) (l^2 - m_2^2) (k + l + p)^2} \\ S^{(2)} &= \epsilon^2 e^{2\gamma_E \epsilon} \int \frac{d^{2-2\epsilon} k}{i\pi^{1-\epsilon}} \int \frac{d^{2-2\epsilon} l}{i\pi^{1-\epsilon}} \frac{m_2^2 - (k + p)^2}{(k^2 - m_1^2) (l^2 - m_2^2) (k + l + p)^2} \\ S^{(3)} &= \epsilon^2 e^{2\gamma_E \epsilon} \int \frac{d^{2-2\epsilon} k}{i\pi^{1-\epsilon}} \int \frac{d^{2-2\epsilon} l}{i\pi^{1-\epsilon}} \frac{m_1^2 - (l + p)^2}{(k^2 - m_1^2) (l^2 - m_2^2) (k + l + p)^2} \end{split}$$

4th master integral is the double tadpole.

Expressions involve Appell  $F_4$ .

Basis of cuts: 3 maximal cuts  $\Gamma_{123}^{(i)}$ , and one 2-line cut  $\Gamma_{12}$  that is the max. cut of the double tadpole.

Ruth Britto, Trinity College Dublin

#### Two-mass sunset



(日) (部) (注) (注) (三)

## Double-edged triangle

4 master integrals: 2 in top topology, Appell  $F_4$ 2 0-mass sunsets in  $p_1^2$ ,  $p_2^2$ 



Ruth Britto, Trinity College Dublin

### Examples of degenerate limits

Take  $p_2^2 \rightarrow 0.$  Basis of master integrals collapses to 2: 1 in top topology, 1 sunset.

Need to construct new dual integration contours.



Obtain coaction by taking the limit, or directly.



Ruth Britto, Trinity College Dublin

Coaction for Feynman Integrals

(日) (部) (注) (注) (三)

## Examples of degenerate limits

In the limit  $p_3^2 \rightarrow 0$ , the integral is reducible to the sunsets. 4-line cuts vanish.



<ロ> <回> <回> < 回> < 回> < 回> < 回> = 三

## **Adjacent Triangles**

$$\begin{split} e^{2\gamma_{E}\epsilon} \epsilon &\left\{ \frac{\Gamma^{2}(1+\epsilon)\Gamma^{4}(1-\epsilon)}{\Gamma(1-2\epsilon)\Gamma(2-2\epsilon)} \frac{p_{1}^{2}-p_{2}^{2}}{p_{2}^{2}} (-p_{1}^{2})^{-2\epsilon} {}_{2}F_{1}\left(1-\epsilon,1-2\epsilon;2-2\epsilon;1-\frac{p_{1}^{2}}{p_{2}^{2}}\right) \\ &-\frac{\Gamma(1+2\epsilon)\Gamma^{3}(1-\epsilon)}{2(1-2\epsilon)\Gamma(1-3\epsilon)} \left[ \frac{p_{1}^{2}-p_{2}^{2}}{p_{1}^{2}} (-p_{2}^{2})^{-2\epsilon} {}_{3}F_{2}\left(1-\epsilon,1,1-2\epsilon;1+\epsilon,2-2\epsilon;1-\frac{p_{1}^{2}}{p_{1}^{2}}\right) \\ &+\frac{p_{1}^{2}-p_{2}^{2}}{p_{2}^{2}} (-p_{1}^{2})^{-2\epsilon} {}_{3}F_{2}\left(1-\epsilon,1,1-2\epsilon;1+\epsilon,2-2\epsilon;1-\frac{p_{1}^{2}}{p_{2}^{2}}\right) \right] \bigg\} \end{split}$$

6 master integrals.



Ruth Britto, Trinity College Dublin

**Coaction for Feynman Integrals** 

Þ 20 / 22

### 4-point integral: Diagonal Box

$$\int_{p_{1}}^{p_{1}} \int_{p_{2}}^{p_{2}} = -e^{2\gamma_{E}\epsilon} \frac{\epsilon(s+t)}{2(1-2\epsilon)} \frac{\Gamma^{3}(1-\epsilon)\Gamma(1+2\epsilon)}{\Gamma(1-3\epsilon)}$$

$$\left[\frac{t^{-2\epsilon}}{s} {}_{2}F_{1}\left(1-2\epsilon,1-2\epsilon;2-2\epsilon;1+\frac{t}{s}\right)\right]$$

$$+\frac{s^{-2\epsilon}}{t} {}_{2}F_{1}\left(1-2\epsilon,1-2\epsilon;2-2\epsilon;1+\frac{s}{t}\right)\right]$$

3 master integrals, 3 natural cuts.



Ruth Britto, Trinity College Dublin

**Coaction for Feynman Integrals** 

# Summary

- Diagrammatic coaction is conjectured to exist, compatible with
  - local coaction on MPLs, eMPLs, …
  - global coaction on hypergeometric functions

Explicitly known at 1-loop. Beyond 1-loop, we find representations for various examples but lack a precise prediction.

- Coaction of L-loop graph has
  - L-loop master integrals in left entries
  - genuine L-loop cuts in right entries

and exhibits a pairing between these objects.

(日) (部) (注) (注) (三)