

## A Method to Distinguish Quiescent and Dusty Star-forming Galaxies with t-SNE

*Friday 28 May 2021 13:35 (5 minutes)*

Large galaxy surveys have revealed a surprising number of galaxies which have ceased (or quenched) their star-formation, seen as they were when the Universe was only half its current age. However, identifying large, but clean, samples of these “quiescent” galaxies has proven difficult. Their spectral shape, as measured by broad-band photometric measurements, are highly degenerate with dusty star-forming galaxies whose light is attenuated by thick clouds of dust and present as an interloper population. We describe a new technique for identifying pure samples of quiescent galaxies based upon t-distributed stochastic neighbor embedding (t-SNE), an unsupervised machine learning algorithm for dimensionality reduction. This t-SNE selection provides an improvement in both purity and completeness over traditional methods (e.g. UVJ, NUVrJ) and over spectral template fitting. We find that t-SNE outperforms spectral fitting in 63% of trials at distances where large training samples already exist. Remarkably, we find evidence for increased performance when applying the method to samples in the even more distant Universe.

**Author:** WEAVER, John (Cosmic Dawn Center, NBI)

**Co-authors:** Prof. STEINHARDT, Charles (Cosmic Dawn Center); MAXFIELD, Jack (summer student)

**Presenter:** WEAVER, John (Cosmic Dawn Center, NBI)

**Session Classification:** Poster session 2

**Track Classification:** Surveys