

Contribution ID: 67

Type: **Talk**

Electromagnetic Counterparts of Neutron Star Mergers: Signatures of Heavy r-Process Nucleosynthesis

Friday 10 June 2022 11:30 (20 minutes)

It has long since been established that observable actinides in the universe originate from the r-process. In 2017, the electromagnetic counterpart to the gravitational wave detection of two merging neutron stars was observed. From the light curve alone it was possible to characterise two ejecta components: one that contains low-Ye material such as lanthanides and possibly actinides, and a high-Ye component with low lanthanide abundances. The dividing characteristic between the two components is the opacity of the material: lanthanides have a ~100 times higher opacity than iron-group material. The opacity of actinides is expected to be on a similar level as that of the lanthanides, or, possibly, even higher.

To identify specific elements, spectroscopic information is required. However, so far no clear detection of individual lanthanides or actinides has been made in the only observed neutron star merger. A great challenge for spectroscopic modelling of kilonovae is the almost non-existent atomic data currently available for lanthanides and actinides. I will present converged and calibrated atomic structure calculations from Zr to U. I will then use this collection of atomic data to show how we can identify signatures or place constraints on the amount of heavy r-process material synthesized in kilonovae.

Author: FLÖRS, Andreas (MPA Garching)

Presenter: FLÖRS, Andreas (MPA Garching)

Session Classification: Supernovae

Track Classification: Supernovae