

The diurnal path to persistent convective self-aggregation

Wednesday, 5 May 2021 16:00 (1h 45m)

Convective self-aggregation (CSA) has attracted a lot of attention as a possible explanation for large scale tropiccal weather phenomena such as the Madden-Julian oscillation and cyclo-genesis. However, CSA is hampered in the realistic limit of fine model resolution when cold pools—dense air masses beneath thunderstorm clouds—are well-resolved.

Here we mimic the diurnal cycle in cloud-resolving numerical experiments by prescribing a surface temperature oscillation. Our simulations show that the diurnal cycle enables the formation of persistent dry patches closely resembling the early onset of CSA. In fact, the dry-patch formation is accelerated by finer resolutions. We attribute these findings to the highly non-linear dynamics of large ‘combined cold pools’ emerging in symbiosis with mesoscale convective systems. Our results may help connecting CSA paradigm in favor of more realistic simulations.

A preprint of the findings are available at: [arXiv:2104.01132](https://arxiv.org/abs/2104.01132)

Primary authors: JENSEN, Gorm Gruner (Atmospheric Complexity, NBI, Uni. Copenhagen); Dr HÄRTER, Jan O. (Atmospheric Complexity, NBI, Uni. Copenhagen); FIÉVET, Romain (Niels Bohr Institute)

Presenter: JENSEN, Gorm Gruner (Atmospheric Complexity, NBI, Uni. Copenhagen)

Session Classification: Modelling and Parameterising Deep Convective Organisation

Track Classification: Modelling and Parameterising Deep Convective Organisation