

## How weakened cold pools facilitate convective self-aggregation

*Thursday, 6 May 2021 16:00 (1h 45m)*

In radiative-convective equilibrium (RCE) simulations, convective self-aggregation (CSA) is the spontaneous organization into segregated cloudy and cloud-free regions. Evidence exists for how CSA is stabilized, but how it arises favorably on large domains is not settled. Using large-eddy simulations (LES), we link the spatial organization emerging from the interaction of cold pools (CPs) to CSA. We systematically weaken simulated rain evaporation to reduce maximal CP radii,  $R_{\max}$ , and find reducing  $R_{\max}$  causes CSA to occur earlier. We further identify a typical rain cell generation time and a minimum radius,  $R_{\min}$ , around a given rain cell, within which the formation of subsequent rain cells is suppressed. Incorporating  $R_{\min}$  and  $R_{\max}$ , we propose a toy model that captures how CSA arises earlier on large domains: when two CPs of radii  $r_{i,j} \in [R_{\min}, R_{\max}]$  collide, they form a new convective event. These findings imply that interactions between CPs may explain the initial stages of CSA.

**Primary author:** Mr NISSEN, Silas Boye (Niels Bohr Institute)

**Co-author:** HAERTER, Jan (Niels Bohr Institute, Copenhagen University)

**Presenter:** Mr NISSEN, Silas Boye (Niels Bohr Institute)

**Session Classification:** RCE and Processes in Deep Convective Organization

**Track Classification:** RCE and Processes in Deep Convective Organization