

Applications of Scattering Amplitudes to Gravitational Waves

Bohr Centennial Meeting

June 21, 2022

Zvi Bern

ZB, C. Cheung, R. Roiban, C. H. Shen, M. Solon, M. Zeng,
arXiv:1901.04424 and arXiv:1908.01493.

ZB, A. Luna, R. Roiban, C. H. Shen, M. Zeng,
arXiv:2005.03071


ZB, J. Parra-Martinez, R. Roiban, M. Ruf. C.-H. Shen,
M. Solon, M. Zeng, arXiv:2101.07254; arXiv:2112.10750

ZB, Kosmopoulos, Luna, Roiban, Teng, arXiv: 2203.06202

UCLA

Mani L. Bhaumik
Institute for Theoretical Physics

Neils Bohr




- **It is really a great honor speaking at conference dedicated to the legacy of Niels Bohr.**
- **What I find marvelous about Niels Bohr is not just the physics, but the way his legacy lives on through the Institute.**
- **The story of what I'm going to tell you about began here at the NBI in 1987, when I met David Kosower. We believed we could find better ways to think about scattering processes.**

Outline

- 1) **Modern approach to perturbative gravity.**
- 2) **Scattering amplitudes as a powerful new tool for gravitational-wave physics.**
- 3) **New results from past 6 months:**
 - **$O(G^4)$ conservative contributions to 2 body interactions.**
 - **A conjecture for determining Kerr Black hole interactions.**
 - **Appearance of new interactions between generic spinning bodies, such as neutron stars.**
- 4) **Issues to be resolved and outlook.**

Outline

Era of gravitational-wave astronomy has begun.




For an instant brighter in gravitational radiation than all the stars in the visible universe are in EM radiation!

How can amplitudes community, help out with core mission of LIGO/Virgo and future detectors?


Can Scattering Amplitudes Help with Gravitational Waves?

What do quantum scattering amplitudes have to do with classical dynamics of astrophysical objects?

unbounded trajectory



bounded orbit



**gauge theories, QCD, electroweak
quantum field theory**

**General Relativity
classical physics**

Black holes and neutron stars are point particles as far as long-wavelength radiation is concerned.

Iwasaki (1971); Goldberger, Rothstein (2006), Porto; Vaydia, Foffa, Porto, Rothstein, Sturani; Kol; Bjerrum-Bohr, Donoghue, Holstein, Plante, Pierre Vanhove; Neill and Rothstein; Levi, Steinhoff; Vines etc


Will explain that scattering amplitudes well suited to push state-of-the-art perturbative calculations for gravitational-wave physics.

Approach to General Relativity

Our approach does *not* start from usual Einstein Field equations.


$$\cancel{R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi GT_{\mu\nu}}$$

~~geometry~~




Gravitons are spin 2 particles

- Not suited for all problems. Works very well for asymptotically flat space-times in context of perturbation theory.
- Well suited for gravitational-wave physics from compact astrophysical objects



Two Body Problem

From Antelis and Moreno, arXiv:1610.03567



- **Small errors accumulate. Need for high precision.**
- **Input to EOB or other modeling to reliably approach merger.**
- **Two primary inputs: binding energy and frequency shift.**

Buonanno and Damour

Post Newtonian Approximation

For orbital mechanics:

Expand in G and v^2

$$v^2 \sim \frac{GM}{R} \ll 1$$



virial theorem

In center of mass frame:

$$m = m_A + m_B, \quad \nu = \mu/M,$$

$$\mu = m_A m_B / m, \quad P_R = P \cdot \hat{R}$$

$$\frac{H}{\mu} = \frac{P^2}{2} - \frac{Gm}{R} \quad \leftarrow \text{Newton}$$

$$+ \frac{1}{c^2} \left\{ -\frac{P^4}{8} + \frac{3\nu P^4}{8} + \frac{Gm}{R} \left(-\frac{P_R^2 \nu}{2} - \frac{3P^2}{2} - \frac{\nu P^2}{2} \right) + \frac{G^2 m^2}{2R^2} \right\}$$

+ ...

1PN: Einstein, Infeld, Hoffmann;
Droste, Lorentz

Hamiltonian known to 4PN order.

2PN: Ohta, Okamura, Kimura and Hiida.

3PN: Damour, Jaranowski and Schaefer; L. Blanchet and G. Faye.


4PN: Damour, Jaranowski and Schaefer; Foffa (2017), Porto, Rothstein, Sturani (2019).

Which problem to solve?

ZB, Cheung, Roiban, Shen, Solon, Zeng

Some problems for (analytic) theorists:

1. Spin.
2. Finite size effects.
3. New physics effects.
4. Radiation.



→ 5. High orders in perturbation theory. ←

Which problem should we solve?

- Want it to be difficult using standard methods.
- Want it to be of direct importance to LIGO theorists.
- Want it to be in a form that can in principle enter LIGO analysis pipeline

2-body Hamiltonian at 3rd post-Minkowskian order

PN versus PM expansion for conservative two-body dynamics

$$\mathcal{L} = -Mc^2 + \underbrace{\frac{\mu v^2}{2} + \frac{GM\mu}{r}}_{\text{non-spinning compact objects}} + \frac{1}{c^2} [\dots] + \frac{1}{c^4} [\dots] + \dots$$

From Buonanno
Amplitudes 2018

$$E(v) = -\frac{\mu}{2} v^2 + \dots$$

non-spinning compact objects

		0PN	1PN	2PN	3PN	4PN	5PN	...
0PM:	1	v^2	v^4	v^6	v^8	v^{10}	v^{12}	...
1PM:		$1/r$	v^2/r	v^4/r	v^6/r	v^8/r	v^{10}/r	...
2PM:			$1/r^2$	v^2/r^2	v^4/r^2	v^6/r^2	v^8/r^2	...
3PM:				$1/r^3$	v^2/r^3	v^4/r^3	v^6/r^3	...
4PM:					$1/r^4$	v^2/r^4	v^4/r^4	...
...						

(credit: Justin Vines)

$$1 \rightarrow Mc^2, \quad v^2 \rightarrow \frac{v^2}{c^2}, \quad \frac{1}{r} \rightarrow \frac{GM}{rc^2}.$$

current known
PN results

current known
PM results

overlap between
PN & PM results

unknown

- **PM results** (Westfahl 79, Westfahl & Goller 80, Portilla 79-80, Bel et al. 81, Ledvinka et al. 10, Damour 16-17, Guevara 17, Vines 17, Bini & Damour 17-18, Vines in prep)

High-energy gravitational scattering and the general relativistic two-body problem

Thibault Damour*

Institut des Hautes Etudes Scientifiques, 35 route de Chartres, 91440 Bures-sur-Yvette, France

(Dated: October 31, 2017)

A technique for translating the classical scattering function of two gravitationally interacting bodies into a corresponding (effective one-body) Hamiltonian description has been recently introduced

“... and we urge amplitude experts to use their novel techniques to compute the 2-loop scattering amplitude of scalar masses, from which one could deduce the third post-Minkowskian effective one-body Hamiltonian.”

tum gravitationally scattering amplitude of two particles, and we urge amplitude experts to use their novel techniques to compute the 2-loop scattering amplitude of scalar masses, from which one could deduce the third post-Minkowskian effective one-body Hamiltonian

Hard to resist an invitation with this kind of clarity!

The recent observation [1–4] of gravitational wave signals from inspiralling and coalescing binary black holes has been significantly helped, from the theoretical side, by the availability of a large bank of waveform templates, defined [5, 6] within the analytical effective one-body (EOB) formalism [7–11]. The EOB formalism combines, in a suitably resummed format, perturbative, analytical results on the motion and radiation of compact binaries, with some non-perturbative information extracted from numerical simulations of coalescing black-hole binaries (see [12] for a review of perturbative results on binary systems, and [13] for a review of the numerical relativity of binary black holes). Until recently, the perturbative results used to define the EOB conservative dynamics were mostly based on the post-Newtonian (PN) approach to the general relativistic two-body interaction. The conservative two-body dynamics was derived, successively, at the second post-Newtonian (2PN) [14, 15], third post-

ntly introduced to derive from the (gauge-invariant) *scattering function* Φ linking (half) the center of mass (c.m.) classical gravitational scattering angle χ to the total energy, $E_{\text{real}} \equiv \sqrt{s}$, and the total angular momentum, J , of the system¹

$$\frac{1}{2}\chi = \Phi(E_{\text{real}}, J; m_1, m_2, G). \quad (1.1)$$

The (dimensionless) scattering function can be expressed as a function of dimensionless ratios, say

$$\frac{1}{2}\chi = \Phi(h, j; \nu), \quad (1.2)$$

where we denoted

$$h \equiv \frac{E_{\text{real}}}{M}; \quad j \equiv \frac{J}{Gm_1m_2} = \frac{J}{G\mu M}, \quad (1.3)$$

with

$$M \equiv m_1 + m_2; \quad \mu \equiv \frac{m_1m_2}{M}; \quad \nu \equiv \frac{\mu}{M} = \frac{m_1m_2}{(m_1+m_2)^2}.$$

0.10599v1 [gr-qc] 29 Oct 2017

Importing QFT Methods

In quantum field theory we are very good at perturbation theory.
Vast experience with gauge theory.

Gravity \sim (gauge theory)²

Unified framework for both gravity and gauge theory.

See Clifford Cheung's talk

Highlight of imported field theory methods:

- 1. Unitarity method for building integrands.** ZB, Dunbar, Dixon, Kosower
- 2. Method of regions.** Beneke and Smirnov
- 3. NRQCD and EFT methods.** Caswell and Lepage; Luke, Manohar, Rothstein
- 4. IBP reductions for Feynman integrals.** Chetyrkin, Tkachov; Laporta
- 5. Method of differential equations for Feynman integrals.**
Kotikov, ZB, Dixon and Kosower; Gehrmann, Remiddi; Henn, Smirnov

Leverage advances in perturbative QCD to help with gravitational waves


Generalized Unitarity Method

Use simpler tree amplitudes to build higher-order (loop) amplitudes.


$$E^2 = \vec{p}^2 + m^2 \leftarrow \text{on-shell}$$

ZB, Dixon, Dunbar and Kosower (1994)

Two-particle cut:




Three-particle cut:



- Systematic assembly of complete loop amplitudes from tree amplitudes.
- Works for any number of particles or loops.

Generalized unitarity as a practical tool for loops.



ZB, Dixon and Kosower;
ZB, Morgan;
Britto, Cachazo, Feng;
Ossala, Pittau, Papadopoulos;
Ellis, Kunszt, Melnikov;
Forde; Badger;
ZB, Carrasco, Johansson, Kosower
and many others


**Idea used in the “NLO revolution” in QCD collider physics.
No gauge fixing in the formalism.**

Simplicity of Gravity Scattering Amplitudes

People were looking at perturbative gravity amplitudes the wrong way.

On-shell three vertices contains all information: $E_i^2 - \vec{p}_i^2 = 0$


QCD:



color factor

$$-gf^{abc}(\eta_{\mu\nu}(k_1 - k_2)_\rho + \text{cyclic})$$

Einstein gravity:



$$i\kappa(\eta_{\mu\nu}(k_1 - k_2)_\rho + \text{cyclic}) \\ \times (\eta_{\alpha\beta}(k_1 - k_2)_\gamma + \text{cyclic})$$

“square” of
Yang-Mills
vertex.

Starting from this on-shell vertex any multi-loop amplitude can be constructed via modern methods

$$\text{Gravity} \sim (\text{gauge theory})^2$$

KLT Relation Between Gravity and Gauge Theory


KLT (1985)

Kawai-Lewellen-Tye string relations in low-energy limit:

$$M_4^{\text{tree}}(1, 2, 3, 4) = -is_{12}A_4^{\text{tree}}(1, 2, 3, 4) A_4^{\text{tree}}(1, 2, 4, 3),$$

$$M_5^{\text{tree}}(1, 2, 3, 4, 5) = is_{12}s_{34}A_5^{\text{tree}}(1, 2, 3, 4, 5) A_5^{\text{tree}}(2, 1, 4, 3, 5) \\ + is_{13}s_{24}A_5^{\text{tree}}(1, 3, 2, 4, 5) A_5^{\text{tree}}(3, 1, 4, 2, 5)$$

Inherently gauge invariant!



Generalizes to explicit all-leg form.

ZB, Dixon, Perelstein, Rozowsky

1. Gravity amplitudes derivable from gauge theory.
2. Once gauge-theory amplitude is simplified, so is gravity.
3. Standard Lagrangian methods offer no hint why this is possible.


Duality Between Color and Kinematics

Zhu; Goebel, Halzen, Leveille

ZB, Carrasco, Johansson (BCJ)


coupling constant \rightarrow color factor \rightarrow momentum dependent kinematic factor

$$-g f^{abc} (\eta_{\mu\nu} (k_1 - k_2)_\rho + \text{cyclic})$$



Color factors based on a Lie algebra: $[T^a, T^b] = i f^{abc} T^c$

Jacobi Identity $f^{a_1 a_2 b} f^{b a_4 a_3} + f^{a_4 a_2 b} f^{b a_3 a_1} + f^{a_4 a_1 b} f^{b a_2 a_3} = 0$



Use $1 = s/s = t/t = u/u$
to assign 4-point diagram
to others.

$$s = (k_1 + k_2)^2 \quad t = (k_1 + k_4)^2$$

$$u = (k_1 + k_3)^2$$

$$\mathcal{A}_4^{\text{tree}} = g^2 \left(\frac{n_s c_s}{s} + \frac{n_t c_t}{t} + \frac{n_u c_u}{u} \right)$$

Color factors satisfy Jacobi identity:

Numerator factors satisfy similar identity:

$$c_u = c_s - c_t$$

$$n_u = n_s - n_t$$

So far nothing more than some curiosity.


Duality Between Color and Kinematics

Consider five-point tree amplitude:

ZB, Carrasco, Johansson (BCJ)

$$A_5^{\text{tree}} = \sum_{i=1}^{15} \frac{c_i n_i}{\prod_{\alpha_i} p_{\alpha_i}^2}$$

color factor
kinematic numerator factor
Feynman propagators



$$c_1 = f^{a_3 a_4 b} f^{b a_5 c} f^{c a_1 a_2} \quad c_2 = f^{a_3 a_4 b} f^{b a_2 c} f^{c a_1 a_5} \quad c_3 = f^{a_3 a_4 b} f^{b a_1 c} f^{c a_2 a_5}$$

$$n_i \sim k_4 \cdot k_5 k_2 \cdot \varepsilon_1 \varepsilon_2 \cdot \varepsilon_3 \varepsilon_4 \cdot \varepsilon_5 + \dots$$

$$c_1 + c_2 + c_3 = 0 \iff n_1 + n_2 + n_3 = 0$$

Claim: We can always find a rearrangement so color and kinematics satisfy the *same* algebraic constraint equations.

Progress on unraveling relations.

BCJ, Bjerrum-Bohr, Feng, Damgaard, Vanhove, ; Mafra, Stieberger, Schlotterer;
 Tye and Zhang; Feng, Huang, Jia; Chen, Du, Feng; Du, Feng, Fu; Naculich, Nastase, Schnitzer
 O'Connell and Montiero; Bjerrum-Bohr, Damgaard, O'Connell and Montiero; O'Connell, Montiero, White;
 Du, Feng and Teng, Song and Schlotterer, etc.

Higher-Point Gravity and Gauge Theory

ZB, Carrasco, Johansson




gauge theory (QCD): $\mathcal{A}_n^{\text{tree}} = ig^{n-2} \sum_i \frac{c_i n_i}{D_i}$

color factor
kinematic numerator factor
Feynman propagators

$$c_k = c_i - c_j$$

$$n_k = n_i - n_j$$

$$c_i \rightarrow n_i$$



sum over diagrams with only 3 vertices

Einstein gravity: $\mathcal{M}_n^{\text{tree}} = i\kappa^{n-2} \sum_i \frac{n_i^2}{D_i}$

$$n_i \sim k_4 \cdot k_5 k_2 \cdot \varepsilon_1 \varepsilon_2 \cdot \varepsilon_3 \varepsilon_4 \cdot \varepsilon_5 + \dots$$


Gravity and gauge theory kinematic numerators are the same!

Same ideas conjectured to hold at loop level.

Cries out for a unified description of gravity with gauge theory, presumably along the lines of string theory.

Web of Theories Linked by Double Copy

See review from ZB, Carrasco, Chiodaroli, Johanson, Roiban





Double copy is a much more general idea than just for gravity

Double Copy for Classical Solutions

Goal is to formulate gravity solutions directly in terms of gauge theory

Variety of special cases:

- Schwarzschild and Kerr black holes.
- Solutions with cosmological constant.
- Radiation from accelerating black hole.
- Maximally symmetric space times.
- Plane wave background.
- Gravitational radiation.



Monteiro, O'Connell and White;
Luna, Monteiro, O'Connell and White;
Luna, Monteiro, Nicholsen, O'Connell and White;
Ridgway and Wise; Carrillo González, Penco, Trodden;
Adamo, Casali, Mason, Nekovar;
Goldberger and Ridgway; Chen;
Luna, Monteiro, Nicholson, Ochirov;
Bjerrum-Bohr, Donoghue, Vanhove;
O'Connell, Westerberg, White; Kosower, Maybee, O'Connell; Adamo, Casali, Mason, Nekovar
etc

**Still no general understanding.
But plenty of examples.**

Effective Field Theory Approach

ZB, Cheung, Roiban, Shen, Solon, Zeng

Cheung, Rothstein, Solon (2018)

**Amplitudes
community**

**Gravitational
Scattering
Amplitudes**

**Effective
Field Theory
Methods**

**EFT
community**

Kawai, Lewellen, Tye

ZB, Dixon, Dunbar and Kosower

ZB, Dixon, Dunbar, Perelstein, Rozowsky

ZB, Carrasco, Johansson; Etc

Beneke, Smirnov (Method of regions)

Goldberger, Rothstein;

Porto; Neill, Rothstein;

Vaydia, Foffa, Porto, Rothstein, Sturani;

Kol, Smolkin, Levi, Steinhoff, etc.

**Post
Minkowskian
Potentials**

**In a form useful for
bound-state problem**

The EFT directly gives us a two-body Hamiltonian of a form appropriate to enter LIGO analysis pipeline (after importing into EOB or pheno models).

We prefer this method when pushing into new territory.

EFT Approach

No need to re-invent the wheel.

Build EFT from which we can read off potential.

Goldberger and Rothstein

Neill, Rothstein


Cheung, Rothstein, Solon (2018)

$$L_{\text{kin}} = \int_{\mathbf{k}} A^\dagger(-\mathbf{k}) \left(i\partial_t + \sqrt{\mathbf{k}^2 + m_A^2} \right) A(\mathbf{k}) \\ + \int_{\mathbf{k}} B^\dagger(-\mathbf{k}) \left(i\partial_t + \sqrt{\mathbf{k}^2 + m_B^2} \right) B(\mathbf{k})$$

$$L_{\text{int}} = - \int_{\mathbf{k}, \mathbf{k}'} V(\mathbf{k}, \mathbf{k}') A^\dagger(\mathbf{k}') A(\mathbf{k}) B^\dagger(-\mathbf{k}') B(-\mathbf{k})$$

two body potential

A, B scalars
represents spinless
black holes



Match amplitudes of this theory to the full theory in classical limit to extract a potential which can then be directly used for bound state.

The EFT is used to define the potential and 2 body Hamiltonian.
This gives us binding energy.

EFT Matching

full general relativity
(complicated)

Amplitude methods
double copy



tree amplitude

$\hbar \rightarrow 0$

generalized
unitarity



loop integrand

Loop integration
Method of regions



GR loop amplitude

effective theory
(simpler)

build
ansatz



potential

Feynman
diagrams



loop integrand

loop
integration



EFT loop amplitude

identical
physics

=

Roundabout but efficiently determines potential.

Alternative Methods

There are now multiple alternative ways that bypass EFT matching and subtraction of iterations.

- **Calculate physical observables** Kosower, Maybee, O'Connell
- **Eikonal Phase** Amati, Ciafaloni, Veneziano;
Di Vecchia, Heissenberg, Russo, Veneziano
- **Amplitude Action Relation** ZB, Parra-Martinez, Roiban, Ruf,
Shen, Solon, Zeng
- **Exponential representation** Damgaard, Plante, Vanhove
- **Heavy mass field theory** Brandhuber, Chen, Travaglini, Wen
Damgaard, Haddad, Helset
- **World line formalisms** Goldberger, Rothstein; Levi, Steinhoff;
Dlapa, Kälin, Liu, Porto;
Jakobson, Mogul, Plefka, Steinhoff;
Edison, Levi

For pushing into new territory we still prefer EFT.


All are fine. Key issue at high orders is efficient loop integration.

General Relativity: Unitarity + Double Copy

- **Long-range force:** Two matter lines must be separated by on-shell propagators.
- **Classical potential:** 1 matter line per loop is cut (on-shell).


Neill and Rothstein ; Bjerrum-Bohr, Damgaard, Festuccia, Planté, Vanhove; Cheung, Rothstein, Solon

Only independent unitarity cut for 2 PM 2 body Hamiltonian.



**Treat exposed lines on-shell (long range).
Pieces we want are simple!**

Independent generalized unitarity cuts for 3 PM.



Our amplitude tools fit perfectly with extracting pieces we want.




gravity




loops

Generalized Unitarity Cuts



2nd post-Minkowskian order



KLT relations

$$\begin{aligned}
 C_{\text{GR}} &= \sum_{h_5, h_6 = \pm} M_3^{\text{tree}}(3^s, 6^{h_6}, -7^s) M_3^{\text{tree}}(7^s, -5^{h_5}, 2^s) M_4^{\text{tree}}(1^s, 5^{-h_5}, -6^{-h_6}, 4^s) \\
 &= \sum_{h_5, h_6 = \pm} it [A_3^{\text{tree}}(3^s, 6^{h_6}, -7^s) A_3^{\text{tree}}(7^s, -5^{h_5}, 2^s) A_4^{\text{tree}}(1^s, 5^{-h_5}, -6^{-h_6}, 4^s)] \\
 &\quad \times [A_3^{\text{tree}}(3^s, 6^{h_6}, -7^s) A_3^{\text{tree}}(7^s, -5^{h_5}, 2^s) A_4^{\text{tree}}(4^s, 5^{-h_5}, -6^{-h_6}, 1^s)]
 \end{aligned}$$

Problem of computing the generalized cuts in gravity is reduced to multiplying and summing gauge-theory tree amplitudes.

$$A_4^{\text{tree}}(1^s, 2^+, 3^+, 4^s) = i \frac{m^2 [23]}{\langle 23 \rangle \tau_{12}} \quad A_4^{\text{tree}}(1^s, 2^+, 3^-, 4^s) = i \frac{\langle 3|1|2 \rangle^2}{s_{23} \tau_{12}} \quad \begin{aligned} \tau_{12} &= 2p_1 \cdot p_2 \\ s_{23} &= (p_1 + p_2)^2 \end{aligned}$$

- **For spinless case, same logic works to all orders: KLT and BCJ work for massless n -point in D -dimension. Dimensional reduction gives massive case.**

Amplitude in Conservative Classical Potential Limit

ZB, Cheung, Roiban, Shen, Solon, Zeng (BCRSSZ)

To make story short. The $O(G^3)$ or 3PM conservative terms are:

$$\mathcal{M}_3 = \frac{\pi G^3 \nu^2 m^4 \log q^2}{6\gamma^2 \xi} \left[3 - 6\nu + 206\nu\sigma - 54\sigma^2 + 108\nu\sigma^2 + 4\nu\sigma^3 - \frac{48\nu(3 + 12\sigma^2 - 4\sigma^4) \operatorname{arcsinh} \sqrt{\frac{\sigma-1}{2}}}{\sqrt{\sigma^2-1}} - \frac{18\nu\gamma(1-2\sigma^2)(1-5\sigma^2)}{(1+\gamma)(1+\sigma)} \right] + \frac{8\pi^3 G^3 \nu^4 m^6}{\gamma^4 \xi} \left[3\gamma(1-2\sigma^2)(1-5\sigma^2)F_1 - 32m^2\nu^2(1-2\sigma^2)^3 F_2 \right]$$

$$m = m_1 + m_2, \quad \mu = m_A m_B / m, \quad \nu = \mu / m, \quad \gamma = E / m, \\ \xi = E_1 E_2 / E^2, \quad E = E_1 + E_2, \quad \sigma = p_1 \cdot p_2 / m_1 m_2,$$

- **Amplitude remarkably compact.**
- **Arcsinh and the appearance of a mass singularity is new and robust feature. Cancels mass singularity of real radiation. No surprise.**
- **IR finite parts of amplitude directly connected to scattering angle.**
- **Derived conservative scattering angle has simple mass dependence.**

Di Vecchia, Heissenberg, Russo, Veneziano; Damour

Expanded on by Kälin, Porto; Bjerrum-Bohr, Cristofoli, Damgaard

Observed by Antonelli, Buonanno, Steinhoff, van de Meent, Vines (1901.07102)

Comprehensive understanding: Damour

Conservative $O(G^3)$ 2-body Hamiltonian

BCRSSZ

The $O(G^3)$ 3PM Hamiltonian: $H(\mathbf{p}, \mathbf{r}) = \sqrt{\mathbf{p}^2 + m_1^2} + \sqrt{\mathbf{p}^2 + m_2^2} + V(\mathbf{p}, \mathbf{r})$

$$V(\mathbf{p}, \mathbf{r}) = \sum_{i=1}^3 c_i(\mathbf{p}^2) \left(\frac{G}{|\mathbf{r}|} \right)^i,$$

Newton in here

$$c_1 = \frac{\nu^2 m^2}{\gamma^2 \xi} (1 - 2\sigma^2), \quad c_2 = \frac{\nu^2 m^3}{\gamma^2 \xi} \left[\frac{3}{4} (1 - 5\sigma^2) - \frac{4\nu\sigma (1 - 2\sigma^2)}{\gamma\xi} - \frac{\nu^2 (1 - \xi) (1 - 2\sigma^2)^2}{2\gamma^3 \xi^2} \right],$$

$$c_3 = \frac{\nu^2 m^4}{\gamma^2 \xi} \left[\frac{1}{12} (3 - 6\nu + 206\nu\sigma - 54\sigma^2 + 108\nu\sigma^2 + 4\nu\sigma^3) - \frac{4\nu (3 + 12\sigma^2 - 4\sigma^4) \operatorname{arcsinh} \sqrt{\frac{\sigma-1}{2}}}{\sqrt{\sigma^2 - 1}} \right. \\ \left. - \frac{3\nu\gamma (1 - 2\sigma^2) (1 - 5\sigma^2)}{2(1 + \gamma)(1 + \sigma)} - \frac{3\nu\sigma (7 - 20\sigma^2)}{2\gamma\xi} - \frac{\nu^2 (3 + 8\gamma - 3\xi - 15\sigma^2 - 80\gamma\sigma^2 + 15\xi\sigma^2) (1 - 2\sigma^2)}{4\gamma^3 \xi^2} \right. \\ \left. + \frac{2\nu^3 (3 - 4\xi)\sigma (1 - 2\sigma^2)^2}{\gamma^4 \xi^3} + \frac{\nu^4 (1 - 2\xi) (1 - 2\sigma^2)^3}{2\gamma^6 \xi^4} \right],$$

$$m = m_1 + m_2, \quad \mu = m_A m_B / m, \quad \nu = \mu / m, \quad \gamma = E / m, \\ \xi = E_1 E_2 / E^2, \quad E = E_1 + E_2, \quad \sigma = \mathbf{p}_1 \cdot \mathbf{p}_2 / m_1 m_2,$$

- Expanding in velocity gives infinite sequence of terms in PN expansion.
- Can be put into EOB form. Antonelli, Buonanno, Steinhoff, van de Meent, Vines

How do we know it is right?

Original check:

Compared to 4PN Hamiltonians after canonical transformation

Damour, Jaranowski, Schäfer; Bernard, Blanchet, Bohé, Faye, Marsat

Thibault Damour seriously questioned correctness.

Specific corrections proposed. [Damour, arXiv:1912.02139v1](#)

Subsequent calculations confirm our 3PM result:

1. Papers confirming our result in 6PN overlap.

Blümlein, Maier, Marquard, Schäfer;
Bini, Damour, Geralico

2. Subsequent calculations reproducing our 3PM result.

Cheung and Solon; Kälin, Liu, Porto

3. Scattering angle checks. [ZB, Ita, Parra-Martinez, Ruf](#)

4. Adding real radiation removes mass singularity.

[Di Vecchia, Heissenberg, Russo, Veneziano; Damour](#)


3PM results have passed highly nontrivial checks and careful scrutiny.

Higher Order Scalability: $O(G^4)$

Methods scale well to higher orders

ZB, Parra-Martinez, Roiban, Ruf,
Shen, Solon, Zeng (2022)


At 4PM or $O(G^4)$ similar except cuts more complicated and integrals significantly harder.



High Loop Integration

ZB, Parra-Martinez, Roiban, Ruf, Shen, Solon, Zeng

Integration more challenging than at 2 loops.



Developed a new hybrid approach that combines ideas from various methods:

1. Method of regions to separate potential and radiation.

Beneke and Smirnov

2. Nonrelativistic integration. Velocity expand and then mechanically integrate. Get first few orders in velocity. Boundary conditions.

Cheung, Rothstein, Solon

3. Integration by parts and differential equations. Imported from QCD.

Chetyrkin, Tkachov; Laporta; Kotikov, Bern, Dixon and Kosower; Gehrmann, Remiddi.

Single scale integrals!

Parra-Martinez, Ruf, Zeng

IBP:
$$0 = \int \prod_i^L \frac{d^D \ell_i}{(2\pi)^D} \frac{\partial}{\partial \ell_i^\mu} \frac{N^\mu(\ell_k, p_M)}{Z_1 \dots Z_n}$$

Solve linear relations between integrals in terms of master integrals.

DEs:
$$\frac{\partial}{\partial s_i} I_j^{\text{master}} = \text{simplified via IBP}$$

Solve DEs either as series or basis of functions.


• Many tools available: We use FIRE6, which is more than sufficient at 3 loops.

• Elliptic integrals make an appearance. At end just a minor annoyance.

Smirnov, Chuharev

Complete Conservative Contribution $O(G^4)$

ZB, Parra-Martinez, Roiban, Ruf, Shen, Solon, Zeng (2022)



test particle 1st self force Iteration. No need to compute

$O(G^4)$ amplitude

$$\mathcal{M}_4^{\text{cons}} = G^4 M^7 \nu^2 |\mathbf{q}| \pi^2 \left[\mathcal{M}_4^{\text{p}} + \nu \left(4\mathcal{M}_4^{\text{t}} \log\left(\frac{p_\infty}{2}\right) + \mathcal{M}_4^{\pi^2} + \mathcal{M}_4^{\text{rem}} \right) \right] + \int_{\ell} \frac{\tilde{I}_{r,1}^4}{Z_1 Z_2 Z_3} + \int_{\ell} \frac{\tilde{I}_{r,1}^2 \tilde{I}_{r,2}}{Z_1 Z_2} + \int_{\ell} \frac{\tilde{I}_{r,1} \tilde{I}_{r,3}}{Z_1} + \int_{\ell} \frac{\tilde{I}_{r,2}^2}{Z_1}$$

$$D = 4 - 2\epsilon$$

$$\mathcal{M}_4^{\text{t}} = r_1 + r_2 \log\left(\frac{\sigma+1}{2}\right) + r_3 \frac{\text{arccosh}(\sigma)}{\sqrt{\sigma^2-1}},$$

$$\mathcal{M}_4^{\text{p}} = -\frac{35(1-18\sigma^2+33\sigma^4)}{8(\sigma^2-1)}$$

tail effect

← elliptic

$$\mathcal{M}_4^{\pi^2} = r_4 \pi^2 + r_5 K\left(\frac{\sigma-1}{\sigma+1}\right) E\left(\frac{\sigma-1}{\sigma+1}\right) + r_6 K^2\left(\frac{\sigma-1}{\sigma+1}\right) + r_7 E^2\left(\frac{\sigma-1}{\sigma+1}\right),$$

$$\mathcal{M}_4^{\text{rem}} = r_8 + r_9 \log\left(\frac{\sigma+1}{2}\right) + r_{10} \frac{\text{arccosh}(\sigma)}{\sqrt{\sigma^2-1}} + r_{11} \log(\sigma) + r_{12} \log^2\left(\frac{\sigma+1}{2}\right) + r_{13} \frac{\text{arccosh}(\sigma)}{\sqrt{\sigma^2-1}} \log\left(\frac{\sigma+1}{2}\right) + r_{14} \frac{\text{arccosh}^2(\sigma)}{\sigma^2-1} + r_{15} \text{Li}_2\left(\frac{1-\sigma}{2}\right) + r_{16} \text{Li}_2\left(\frac{1-\sigma}{1+\sigma}\right) + r_{17} \frac{1}{\sqrt{\sigma^2-1}} \left[\text{Li}_2\left(-\sqrt{\frac{\sigma-1}{\sigma+1}}\right) - \text{Li}_2\left(\sqrt{\frac{\sigma-1}{\sigma+1}}\right) \right].$$

$$\nu = m_1 m_2 / (m_1 + m_2)^2$$

$$\sigma = p_1 \cdot p_2 / m_1 m_2,$$

r_i rational coefficients

This is complete conservative contribution.

$$\mathcal{M}_4^{\text{radgrav,f}} = \frac{12044}{75} p_\infty^2 + \frac{212077}{3675} p_\infty^4 + \frac{115917979}{793800} p_\infty^6 - \frac{9823091209}{76839840} p_\infty^8 + \frac{115240251793703}{1038874636800} p_\infty^{10} + \dots$$

First 3 terms match 6PN results of Bini, Damour, Geralico!

- Some disagreement in one term with 6PN of Blumlein, Maier, Marquard, Schafer.
- Claim by Diapa, Kalin, Liu, Porto that we dropped “memory” pieces. *We disagree.*


Issues to be Resolved

- 1) Disagreements with Bluemlein et al and especially Porto et al still needs to be fully resolved. Recent progress.

Damour's "Good Polynomiality": No $1/(m_1 + m_2)$. Manifest in our scattering amplitude formalism.

Forthcoming paper from Bjerrum-Bohr, Vanhove, Plante, confirms our result.

- 2) We give a Hamiltonian valid for (unbound) hyperbolic motion. But analytic continuation (bound) elliptic motion at $O(G^4)$ is nontrivial. This issue is identical to the one encountered in PN approximation. Needs to be resolved.




Comparison with Numerical Relativity


Khalil, Buonanno, Steinhoff, Vines

As an interesting check compare to numerical relativity:

Original angle in PM perturbation



EOB-improved angle



Numerical data from Damour, Guercilena, Hinder, Hopper, Nagar, Rezzolla

Surprisingly good agreement with numerical relativity.

Dissipative Contribution to Impulse

The impulse or angle also depends on dissipative effects.

(usually separated for bound states).

Manohar, Shen and Ridgeway computed odd terms in velocity at $O(G^4)$.

Di Vecchia, Heissenberg, Russo, Veneziano;
Damour; Herrmann, Parra-Martinez, Ruf, Zeng
Manohar, Shen and Ridgeway;
Bjerrum-Bohr, Vanhove, Plante (to appear)

See talks from Di Vecchia
Veneziano

$$\Delta p_{\perp,4} = \nu M^5 \left(\frac{G}{b} \right)^4 (c_{b,4}^{\text{cons}} + c_{b,4}^{\text{rr,even}} + c_{b,4}^{\text{rr,odd}})$$

$$c_{b,4}^{\text{rr,odd}} = \nu \left[\frac{\sigma(6\sigma^2 - 5)}{\sigma^2 - 1} - \frac{m_1}{M} \frac{2\sigma^2 - 1}{(\sigma + 1)} \right] \frac{\mathcal{E}(\sigma)}{p_{\infty}} - \frac{\nu(2\sigma^2 - 1)}{\sigma^2 - 1} \left[\frac{3\pi(5\sigma^2 - 1)}{2} \mathcal{I}(\sigma) + \mathcal{C}(\sigma) + 2\mathcal{D}(\sigma) \right]$$

$$\mathcal{I}(\sigma) = -\frac{16}{3} + \frac{2\sigma^2}{\sigma^2 - 1} + \frac{4(2\sigma^2 - 3)}{\sigma^2 - 1} \frac{\sigma \operatorname{arcsinh} \left(\sqrt{\frac{\sigma-1}{2}} \right)}{\sqrt{\sigma^2 - 1}}$$

$$\frac{\mathcal{E}(\sigma)}{\pi} = f_1 + f_2 \log \left(\frac{\sigma + 1}{2} \right) + f_3 \frac{\sigma \operatorname{arcsinh} \left(\sqrt{\frac{\sigma-1}{2}} \right)}{\sqrt{\sigma^2 - 1}}$$

$$\frac{\mathcal{C}(\sigma)}{\pi} = g_1 + g_2 \log \left(\frac{\sigma + 1}{2} \right) + g_3 \frac{\sigma \operatorname{arcsinh} \left(\sqrt{\frac{\sigma-1}{2}} \right)}{\sqrt{\sigma^2 - 1}}$$

$$\mathcal{D}(\sigma) = \frac{3\pi(5\sigma^2 - 1)}{8} \mathcal{I}(\sigma)$$

f_i and g_i are rational functions.

Energy loss still needed for longitudinal impulse.

4PM is under good control, but still some cleaning up needed.

Outlook

Amplitude methods continue to show a lot of promise.
They have already been tested for a variety of nontrivial problems.

- **Pushing state of the art for high orders in G.**
ZB, Cheung, Roiban, Parra-Martinez, Ruf, Shen, Solon, Zeng
- **Radiation.** Cristofoli, Gonzo, Kosower, O'Connell; Herrmann, Parra-Martinez, Ruf, Zeng; Di Vecchia, Heissenberg, Russo, Veneziano
- **Finite size effects.** Cheung and Solon; Haddad and Helset; Kälin, Liu, Porto; Cheung, Shah, Solon; ZB, Parra-Martinez, Roiban, Sawyer, Shen
- **Spin.** Vaidya; Geuvara, O'Connell, Vines; Chung, Huang, Kim, Lee; ZB, Luna, Roiban, Shen, Zeng; Kosmopoulos, Luna; Damgaard, Haddad, Helset; Aoude, Haddad, Helset

↑
We have more Wilson coefficients than our GR friends. Stay tuned...

ZB, Kosmopoulos, Luna, Roiban, Teng

Ask Andres Luna

Amplitude methods have become a standard tool for gravitational-wave problems.

Summary

- Scattering amplitudes give us new ways to think about problems of current interest in general relativity.
- Double-copy idea gives a *unified framework* for gravity and gauge theory.
- Combining with EFT methods gives a powerful tool for gravitational-wave physics in language LIGO/Virgo can use.
- Pushed state of the art: $O(G^4)$ Newtonian-like conservative part.
- Methods nowhere close to exhausted.
- Higher orders in G , resummations in G , spin, finite-size effects, radiation and dissipation obvious paths to pursue.

Tools developed over the years for carrying out perturbative QCD computations are well suited for gravitational-wave problem and are being used to push the state of the art.