



Exercise 1: Interactions of relativistic protons

Maria Petropoulou

July 2025

1. The relation expressing the energy threshold condition for the two-particle interaction $a + b \rightarrow c + d$ is:

$$E_a E_b - \vec{p}_a \vec{p}_b c^2 \geq \Delta m c^4 (m_a + m_b + \Delta m/2) + m_a m_b c^4 \quad (1)$$

where $\Delta m = m_c + m_d - (m_a + m_b)$, and m_i, E_i, \vec{p}_i are the rest mass, total energy, and momentum, respectively, of the particle i .

- (a) Calculate the energy threshold for the interaction $p + \gamma \rightarrow e^- + e^+$. (Consider that the photon target has a fixed energy E_γ in the lab frame).
 - (b) Calculate the energy threshold for the interaction $\frac{A}{Z}X + \gamma \rightarrow \frac{A}{Z}X + \pi^0$. (Consider that the photon target has a fixed energy E_γ in the lab frame).
 - (c) An ultra-high-energy cosmic ray (UHECR) proton and an iron nucleus are traveling through the Universe and are interacting with photons from the cosmic microwave background (CMB) via the photopair process. Using the expression you derived above, calculate the minimum energy of the UHECR proton/iron nucleus for a head-on collision with the CMB photons. Comment on your results. [$m_e c^2 \approx 0.5$ MeV, $m_p c^2 \approx m_n c^2 \approx 1$ GeV, $m_{\pi^0} c^2 \approx 135$ MeV, $\epsilon_{CMB} \approx 6 \times 10^{-4}$ eV].
2. The mean free path for energy losses of an UHECR proton with Lorentz factor γ_p due to $p\gamma$ interactions with CMB photons is given by:

$$\frac{1}{\ell(\gamma_p)} = \frac{1}{2\gamma_p^2} \int_{\bar{\epsilon}_{th}}^{\infty} d\epsilon \bar{\epsilon} \hat{\sigma}(\bar{\epsilon}) \int_{\bar{\epsilon}/(2\gamma_p)}^{\infty} d\epsilon' \frac{n_{CMB}(\epsilon')}{\epsilon'^2} \quad (2)$$

where $\hat{\sigma} = 70 H(\bar{\epsilon} - \bar{\epsilon}_{th}) \mu b$ is an approximate expression for the effective cross section, $\bar{\epsilon}_{th} \simeq 400$ and $H(x)$ is the Heaviside function. All energies in the expression above are normalized to $m_e c^2$.

- (a) The differential CMB photon number density can be approximated as a power law with a sharp cutoff: $n_{CMB}(\epsilon) = n_0 \epsilon$, for $\epsilon \leq \epsilon_{CMB}$, where $\epsilon_{CMB} \simeq 6 \times 10^{-4}$ eV. Calculate




Figure 1: Mean free path for $p\gamma$ interactions of UHECR protons with ambient photon fields.

the normalization n_0 if you know that the local CMB energy density is $u_{CMB} \approx 0.25 \text{ eV cm}^{-3}$.

- (b) Make a plot of $\bar{\ell}(E_p)$ versus E_p (*Hint*: think about the upper limits of integration). How does the approximate solution found in the previous question compare with the accurate result shown in Fig. 1? Comment on your results.