

Lectures: on "beyond" linear GR on "beyond" GR

Refs:

(1A) • Damour 1982 "Surface effects in BH physics"

• arxiv: 1101.2451 NS $\rightarrow$ Einstein

• " : 1202.05286 Gal / Carroll

• " : 1903.09654 Carroll Physics

• " : 2212.06175 Non Linear BH...

• " : 2310.15427 F-G & BDNK

To speak of non-linear science is like
referring to the bulk of biology as the study
of non-elephant animals (S. Ulam)

Context What to expect in observables?
perturbative approaches


- . regime assumptions
- First pples understanding
- . Using right intuition?

strategy:

Back told" efforts in a new light

Connect with physics we know better or with different tools


Consider different surfaces in
asympt. dS ($\Lambda < 0$)



Introduce a null foliation,

$$ds^2 = -A du^2 - 2B du dr + r^2 h_{AB} (dx^A - U^A du) (dx^B - U^B du)$$





- Note: 1) since $u = \text{const}$ is null
this is a "more rigid" choice!
for coords

2) Einstein Eqs have a
particular structure.

•) $E_{AB} - \frac{1}{2} g_{AB} E_{CD} g^{CD} = 0 \Rightarrow \dot{h}' \sim h''$

•) $E^u_a = 0 \quad \left\{ \begin{array}{l} A_{,n} \dots U^A_{,rr} \dots \\ B_{,n} \dots \end{array} \right.$

•) $E^r_a \rightarrow$ constraints that
are propagated
by Bianchi ids.

Aside in a Cauchy foliation...

$\int_{\Sigma} n^a$ $E_{ab} n^a n^b \rightarrow \text{HAMILT. CONST}$

$$\sim E_{ab} n^a \frac{1}{c} \rightarrow \text{Mom Const.}$$

$$E_{ab} \frac{1}{c} \frac{1}{c} \rightarrow (\gamma_{cd}^{ab})_{ct} \gamma_{et}^{ab} \\ \text{(very roughly).}$$

$\Rightarrow$ Bianchi identities $\Rightarrow$

$$\text{if } \{H=0, M_i=0\}_{t=0}$$

$$\text{Evol Eqs} \Rightarrow \dot{H} \sim F_H(H, M, EE) \\ \dot{M}_i \sim F_{M_i}(H, M, EE)$$

In the null foliation.

$$\rightarrow \text{if } \{R^r_a=0\}_{at} \text{ at } r=R$$

$\rightarrow$ propagated through

Aside 2 Relativistic Hydrodynamics
& 'c limits'

Starting point: stress-energy tensor

$$T^{ab} = (\epsilon + p) \frac{u^a u^b}{c^2} + p g^{ab} + \tilde{\tau}^{ab} + \frac{u^a q^b + u^b q^a}{c^2}$$

ϵ : energy density

p : pressure

u^a : fluid velocity

$$\nabla_a T^{ab} = 0$$

$$u^a u_a = -c^2$$


Now, let's take limits:

Galilean limit $c \rightarrow \infty$

$$- u^0 = \gamma c = c + \dots$$

$$- u^i = \gamma v^i = v^i$$

$$- \epsilon = (e + c^2) \rho_0$$



"open the cone"

Now, how about the equations?

$$- c \nabla_\mu T^{\mu 0} = c^2 A + \epsilon + \mathcal{O}(1/c^2)$$

$$- \nabla_a T^a_i = M_i + \mathcal{O}(1/c^2)$$

$A=0 \rightarrow$ continuity equation

$E=0 \rightarrow$ energy conservation

$M_i=0 \rightarrow$ mom. conservation

Take for simplicity, flat spacetime

- $C \approx \partial_t \phi + \partial_i (\phi v^i)$

- $E \approx \partial_t e + v^i \partial_i e \dots$

- $M_i \approx \partial_t (\phi v_i) + \partial_j (\phi v^j v_i) + \dots$


$\Rightarrow$ Navier-Stokes eqn

$\leadsto$ Galilean limit ($C \rightarrow \infty$)

Carrollian limit $C \rightarrow \infty$

$v \rightarrow 0$ faster than $C \rightarrow \infty \rightarrow v^i = C^2 \beta^i$

$\leadsto \begin{cases} u^0 \approx C \\ u^i \approx C^2 \beta^i \end{cases}$



$\rightarrow \begin{cases} C \nabla_a T^a{}_c = \frac{1}{C^2} \tilde{T} + \mathcal{E} + \mathcal{O}(C^4) \\ \nabla_a T^{ai} = \frac{1}{C} H^i + G^i + \mathcal{O}(C^2) \end{cases}$

→ Doubling of equations?

no in the perfect fluid case $F=H=0$
end

$$\mathcal{E} = -\partial_t \epsilon - (\epsilon + p) \Theta$$

$$G_T = (\epsilon + p) \left[\psi_T + \delta_T^c + \beta_T \Theta^c \right] \\ + \beta_T \partial_t (\epsilon + p) + \hat{\partial}_T P$$

~~~~~

Some qns.

1. How does a C. fluid  
equilibrate if the  
cones are closed



2. Causal symmetries

2 - Can't make symmetries,  
where else?

3 - What is a C. eqn  
of state?

4 - What is such a fluid?

Why do we care?

- I care about gravity! Fluids are  
a different class of beasts!

- linearly deg. vs truly non linear!

•  $u_t = u_x$  vs  $u_t = u u_x$

smooth solns > shocks!

- smooth joins  
from smooth data

- smooths:  
from smooth data

- prop. speeds

$c$  or gauge  
order. of state  
of field


- dependent of  
state of field  
 $v = c_s + v_0$

- Why would we learn anything  
useful?!

- "only" non-linear classical field  
that treats non-linearities seriously...

Fluid-gravity correspondence...

[Damour] What do we get if  
one projects  $EE_S$  onto  $\mathcal{H}$ ?



- define  $T_A = - n_a l^a_{jA}$

$\partial_{\mu} T^{AB} \propto \partial_{\mu} T^{AB}$

$$n^a l_a = 1$$

$\gamma_{AB}$  int. metric  
of "Cut"

$$\sigma_{AB} = \frac{1}{2} \frac{D \gamma_{AB}}{dt} - \frac{1}{2} \theta \gamma_{AB}$$

- Raychaudhuri eqn.

$$\dot{\theta} - \kappa \theta + \frac{1}{2} \theta^2 + \sigma_{AB} \sigma^{AB} = 0$$

- Damour (-NS) eqn

$$\dot{\pi}_A + \theta \pi - \nabla_A \kappa - \frac{1}{2} \nabla_A \theta + \nabla^B \sigma_{AB} = 0$$

With the identifications:

$$\epsilon = \theta, \quad \rho = -\kappa \rightarrow \text{EOS!}$$

$$\pi_{AB} = 2\eta \sigma_{AB} + \xi \theta \gamma_{AB}; \quad \eta = \frac{1}{2}, \quad \xi = -\frac{1}{2}$$

$$\pi_A = -\frac{1}{\kappa} \left( \nabla_A \kappa - H^B \dot{\gamma}_{AB} - \frac{1}{\kappa} H^A \dot{\kappa} \right)$$

$$\Sigma^{AB} = J^A = 0$$

$\Rightarrow$  Collisional hydrodynamics

$\rightarrow$  for a particulate fluid.

From gravity  $\rightarrow$  fluids

- How does a fluid with "closed" cones thermalizes?

From fluid  $\rightarrow$  gravity

- How about non linear behavior?

For Collision fluids dual to horizons we can link these questions!

[Redmond-Yorke-LL]

- But, how to close the system of eqs?

- $k$ ?,  $\sigma_{AB}$ ?

$\rightarrow$  dissipation/heating?

$\rightarrow$  EOS?

$\rightarrow k, \sigma_{AB}$  are "sensitive" to the bulk! null surface is  $C$  d+2 s.t.:

→ Involve rest of EES!

$R_{AB}$

$$\bullet \Theta^{(1)} + 12 \Theta^{(2)} + \Theta^{(4)} \Theta^{(6)} - [D_A H^A + H_A H^A] + \frac{R}{2} = 0$$

$$\bullet 2 \sigma_{AB}^{(4)} - 4 \sigma_{(A}^{(2)} \sigma_{B)} + \Theta^2 \sigma_{AB} + (2K - 8) \sigma_{AB}^{(2)} + R_{AB}^{TF} - 2(H_A H_B)^{TF} - 2[D_A H_B]^{TF} = 0$$

— Horizon is teleological!

$$\Theta \rightarrow 0$$

$$u \rightarrow \infty$$

→ In Carrollian picture

$\Theta \rightarrow$  energy and "cup"

• fluid thermalizes by "deforming" the cup!

• Horizon "uniformizes" by adjusting behavior of generators!

Back to exercise!



perturbatively

- perturbatively & resummation

$$\begin{aligned} -\dot{H}_A &= \nabla^2 H_A + \nabla_A \nabla_B H^B \\ &+ \frac{1}{m} [H_A \nabla^A H_B + H_B \nabla^B H_A] \\ &+ \dots \end{aligned}$$

Reynolds number.

$$\nabla^2 H_A + \nabla_A \nabla_B H^B \ll H \nabla H ?$$

$\Rightarrow$  condition

+ non-linear mode generation, etc...

Let's go to  $r \rightarrow \infty$  [AdS bdy]

- consider a perturbed BH

$$\Rightarrow \text{typically } g = g_B + h_1 + h_2 + \dots$$


Instead consider

$$g = g_B(M, a) \rightarrow g(M(x), a(x))$$

and assume a gradient expansion

$$\text{i.e. } |\partial^n F| > |\partial^{n+1} F|$$

- Plug this into EEs or Characteristic form e.e.



$R_{ra} \rightarrow$  "constraints on  $\tau^a$ "

Rest Radial eqs inwards

$$\rightarrow D_a(T^{ab}) = 0 \rightarrow \text{for a symmetric tensor}$$

$$\rightarrow \nabla^a_a = 0$$

Say the BH was a "boosted" black brane with boost  $u^a$ , radius  $r_H = \frac{1}{b}$

$$T^{ab} = \frac{3}{b^4} u^a u^b + \frac{1}{b^4} (u^a u^b + \gamma^{ab})$$

$$P = \frac{\epsilon}{3} \quad (\text{trace condition})$$

1<sup>st</sup> order in gradients?

$$T_{ab} = \frac{3(1+\chi)}{b^4} + \frac{(1+\chi)}{b^4} (u^a u^b + \eta^{ab}) \\ - \frac{2}{b^3} \sigma_{ab} - \frac{8}{b^4} (\eta_a \eta_b)$$

with  $\sigma_{ab} = P^{cd} \partial_c u^b - \frac{1}{3} P^{ab} \partial_c u^c$

$$P^{ab} = u^a u^b + \eta^{ab}$$

$\chi, \eta^a$  = "corrections" which  
are gauge dependent.

- Notice

- transport coeff for shear vis.  
defined

- Can choose  $\chi = \eta^a = 0$

[Bhattacharyya  
'2008].

- Can go to even 2<sup>nd</sup> order  
& 1-to-1 correspondence  
with Landau-Lifschitz!
- Pert's of shear & sound modes  
↔ Lowest QNMs of BH.

What else?

- long-wavelength pert  
→ turbulent regime?  
• turbulence in gravity???

- Freedom in  $X, \alpha$ ?

in AdS

6:16 Sat, Jan 17 GM • **Beyond astro: AdS/CFT...**

92%

- The AdS/CFT correspondence relates a d-dimensional QFT with a (d+1)-dimensional gravity.




[Image: J. Santos]


- Any gravitational phenomena should have an equivalent CFT analog, *and vice versa*.
- A natural arena to study field theory open questions: transport properties in strongly coupled field theories, quantum quenches, thermalization, etc.
- Plenty of applications. Most of which in equilibrium situations and in the probe limit (phase space analysis) (e.g. CMT applications)
- More recently, *dynamical applications*: e.g. examining quark-gluon plasma behavior, thermalization,...
- But also... new physics in gravity?

## Gravitational turbulence?

- Einstein eqns do not develop shocks (propagation speeds = c, regardless of the state of the solution -> no crossing of characteristics). \*VERY\* different from fluids!
- But, AdS/CFT motivates (& makes 'rigorous') a fluid/gravity correspondence, previously identified through a 'membrane paradigm programme' [Damour, Price-Thorne-McDonald] -> [Hubeny-Rangamani-Minwalla-Bhattacharya]
  - Relies on a 'non-standard' [in GR] perturbative approach



[Adams, Chesler, Liu. PRL 2014]



[Green, Carrasco, LL, PRX 2013]



## in AF) Perturbative analysis

- The problem is non-linear....what can we draw from turbulence in AdS?
- *Back to AF spacetimes*, beyond linear perturbation? Take  $h_1$  (with  $l, m, n$ ); a further mode on the 'time-dependent' background, obeys:

$$[ \Box_{\text{kerr}} + O(h_1) ] \Phi = 0.$$

- With the solution having the form:  $e^{t(\alpha - \omega_l)}$  with

$$\alpha = \pm \sqrt{|H h_0(t)/Q m'|^2 - (\omega'_R - \omega_R/2)^2},$$





- So exponentially growing solution if:

$$h_0(t)/(m' \omega'_l) - |Q/H| \sqrt{(\omega'_R - \omega_R/2)^2 / \omega'^2_l + 1} > 0.$$

- if  $\Phi$  has  $l, m/2 \rightarrow$  a parametric instability can turn on; i.e. inverse cascade.
- Further, one can find 'critical values' for growth onset.
- And can define a max value as:
 
$$\text{Re}_g = h_0/(m \omega_v)$$
- identify  $\lambda \leftrightarrow 1/m$  ;  $v \leftrightarrow h_0$  ;  $\eta/\rho \leftrightarrow \omega_v$ 

$$\rightarrow \text{Re}_g = \text{Re}$$

## 'Turbulence', AF spacetimes and 'beyond linear'





$a = 0.998$ , perturbation  $\sim 0.02\%$ , initial mode  $l=2, m2$

Could 'potentially' have observational consequences. Signal is different from what might be expected at the linear level

[Yang, Zimmern, LL '14]




## Exploring non-linearities in FD



# Full non-linear

quasi) 'Steady state' analysis

- 'Stirr' a (region of) spacetime with incoming GWs:  $F(A, \omega, \ell, m=0)$
- Choices :
  - non-spinning BH
  - $\omega = 1/2m ; 1/m ; 2/m$
  - $\ell = 2, 6$
  - $A = \text{'small', 'large' } [R_{\text{eg}}]$
  - Drive with linear growth until  $t_*$ , steady afterwards





Gauge choice?  $\{f^a, \chi\}$

- Unperturbed BH.  $\begin{cases} K = \frac{2}{b}; & \Theta = 0 \\ \pi_a = 0; & \varphi_a = 0 \end{cases}$

- 1<sup>st</sup> order in gradient...

$$\bullet K = \frac{2}{b} - \frac{2}{3}\Theta + \frac{\chi}{2b} + \frac{u^a \partial_a b}{b}$$

$$\bullet \Theta = \Theta - 3 \frac{u^a \partial_a b}{b}$$

$$\bullet \pi_a = \frac{2}{b} f_a - \frac{P_a^b \partial_b b}{b}$$

Options?:  $\pi_a = 0 \iff \partial^a \parallel u^a$

$$\hookrightarrow f^a = -b \partial^a + P^{ab} \partial_b b$$

$$\therefore r_H = \frac{1}{b} + \frac{\Theta}{6} + \frac{\chi}{4b}$$

at 1<sup>st</sup> order  $\Theta = 0 \Rightarrow \chi \propto \Theta$  keep  $r_H$



$$\hookrightarrow \chi = \alpha b \oplus$$

Implications?

$$\Rightarrow \mathcal{E}^U = \frac{3}{b^3} \alpha \left( \partial_a u^a + \frac{u^a \partial_a e}{e+p} \right)$$

$$Q^a = \frac{4}{b^3} \left( Q^a + \frac{P^{ab} \partial_b e}{3(e+p)} \right)$$

$\Rightarrow$  Of the type argued by BDNK!

- 1<sup>st</sup> order in gradient Tab
- contains dissipation
- is strongly hyperbolic  $\rightarrow$  well posedness!
- by passes them! (entropy "issue")

QN Why does a gauge "natural"  
 horizon description

for this you description  
"leads" to BDNK?

---

Deeper analysis of GWs

↳ for beyond GR

↳ for beyond "ingredients"

a.k.a. how would one tell?

① Null tests → Model A vs B  
GR vs Beyond GR

⇒ requires understanding well A  
and/or "smoking guns" for B.

⇒ Options

\* Identify a theory  
- predict deviations

\* Agnostic approach

Theory? What theory? (where from)  
- what changes?  
- how to study it?  
(where and how?)

Derive theories in an EFT way...

Ex Euler vs Navier Stokes

incompressible fluid example

$$\underbrace{\partial_t v + v \partial v}_{\text{"bare theory"}} = \underbrace{\left( \begin{smallmatrix} \eta \\ \zeta \\ \rho \end{smallmatrix} \right) \partial^2 v}_{\text{"correction"}}$$

- Notice: • correction changes character  
→ hyperbolic to parabolic!

For GR ...  $\mathcal{L} \Rightarrow R + \wedge R^p f(\phi)$

•  $R^p$ : curvature comb of order  $p$

- $f(\phi)$ : possible extra d.o.f (beyond those on  $g_{ab}$ )

Eg: ESGB :  $\phi [R^2, R_{ab}R^{ab}, R_{abcd}R^{abcd}]$

dCS :  $\phi R_{abcd} \star R^{abcd}$

Cubic :  $R_{ab}{}^{cd} R_{cd}{}^{lm} R_{lm}{}^{ab}$

Predictions?

Merger (fully non-linear regime in pph)

Eob • ESGB...  $\rightarrow \Box g = \Lambda \partial^2 g \partial^2 g \partial_{ab} g$   
 $\rightarrow$  character can/does change!

• Cubic  $\rightarrow \Box g = \Lambda \partial^3 g$  !

$\rightarrow$  unclear if relevant problems are well posed!

$\rightarrow$  though behaviour in GR helps!

→ if it breaks, EFT doesn't work  
needs something else  
(e.g. collapse scenarios!)

How to assess?

Option 0 :  $\Box g = \lambda S(g^P)$

Red. Ord. → secular behavior kills it!

Option 1 :

Fix eqs

$$\Box g = \lambda \pi$$

$$\Box \pi = k \partial_t \pi + \sigma'(-\pi + S)$$

Option 2 :  $\int (\partial \phi)^2 + \lambda (\partial \phi)^4$

Redef. with  
higher order  
op.

$$\phi \rightarrow \phi + \sigma' \Box \phi$$

$$\hookrightarrow \int (\partial \phi)^2 + \lambda (\partial \phi)^4 + \sigma' (\partial \phi) \Box \phi$$

$$\hookrightarrow \Box \phi = \lambda S + \sigma' \Box \Box \phi$$

Inspired : use pert approaches

(PN, NRGR, amplitudes)

but not a distinction

and compute precociously

Ring down:

- Find solns for BHs
- Compute QNM behaviour.

But in what theory!?

Alternatives?

- parameterize deviations

PPN  $\varphi(\xi) = \sum_i (C_i^{\text{GR}} + \delta_i) f^i$

→ Search for  $\delta_i$  on data

→ can one do better?

Note: evidence so far points to...

- i) GR does quite well!
- ii) If new scale  $L \rightarrow L \lesssim \tilde{L} \sim \text{km}$
- iii) in GR no significant transfer of energy towards UV  
→ EFT!



Recall  $h$  is dimensionless!

$$\leadsto h \simeq h_{GR} + \delta$$

What can one say about  $\delta$ ?

$$\textcircled{1} \quad \delta \sim \left(\frac{L}{M}\right)^q \delta_b$$

$$\textcircled{2} \quad \delta_b \sim \text{lead order PN or inspiral.}$$

Strategy

$$\textcircled{1} \quad \mathcal{R} + \Lambda \ddot{\mathcal{R}}''^p \rightarrow \Lambda = L^{2(p-1)}$$

$$\rightarrow q = 2(p-1)$$

$$\textcircled{2} \quad \mathcal{R} + \Lambda \phi \ddot{\mathcal{R}}''^p \rightarrow \Lambda = L^{2(p-1)}$$

$$\leadsto \text{but } q = 2(2[p-1])$$

EOM

$$\rightarrow G_{ab} \approx \nabla_a \phi \nabla_b \phi \quad (\Lambda^2) \leftarrow$$

$$\Box \phi = \Lambda \phi \quad \rightarrow$$

Can follow this in general for any case!

$\rightarrow q \rightarrow \text{integer}$ ; in several events  
look for  $\delta \sim \frac{1}{M^q}$

$|q| \geq 4$  from EFT in vacuum

$q = 0$  will imply on ECO!

Example:  $q=4 \rightarrow$  ESGB  
dCS  
cubic ✓

• will eliminate  $R_{-} R_{-} R_{-} R_{-}$   
and others  
but one can do better.

EFT "rules": Consider all operators  
consistent with symmetries

and assumptions see...

it can't be ESGB or dCS

→ it must be both!

→ one can then ask which one to compute?

→ guess: circular case ( $\frac{v}{c}$ ) is small

⇒ in inspiral ESGB leads

⇒ if  $\phi$  is present → dipole radn

cubic → only starts at 5PN

**SMOKING GUN**

$q=4$  &  $PN < 5$

→ ESGB + dCS

(and one can go even deeper)

New strategy : multiple events at a time

• search for  $q$  &  $PN$

→  $q \geq 4$  & integer Beyond GR

→  $q=0 \rightarrow \text{ECO!}$

→  $q = 1 \leftrightarrow 2$  environmental effect

→  $q$  other astrophysical



## Final thoughts

- × Useful to think of problems anew
- × In particular to gain new inspiration & tools from other fields

