

Recent ATLAS results:  
SM Higgs Boson Searches

“Spåtind 2012”


Nordic Conference on Particle Physics

A. L. Read (U. Oslo/LAL)



# Outline

- Combination statistics and LEE refresher
- Status of SM Higgs search, November 2011
- Individual ATLAS SM Higgs search results
- Combined ATLAS results, December 2011





# Profile Likelihood ratio, $p_0$ and CLs...in 60 seconds



$$\hat{\mu} \approx \mu_{true} \pm \sigma_\mu, (\mu_{true} = 0, 1, \text{whatever})$$



- $p_0$  to test background hypothesis
- $CL_s$  to test signal hypothesis (e.g.  $\mu=1$ )
- Signal limits on  $\mu$  for  $CL_s = \text{e.g. } 5\%$
- $\hat{\mu}$  to estimate signal strength





[Follow LHCHCG Combination Procedures](#)

# Look-elsewhere effect



- Consider a search with predicted background  $10^{-7}$ 
  - 1 observed event is a  $5.2\sigma$  discovery
- Now consider  $10^7$  searches each with  $10^{-7}$  background
  - Expect on the average 1 event with local p-value of  $10^{-7}$  but this is certainly not a discovery!
  - Probability to make a false discovery is  $P(n \geq 1 | b = 1) = 1 - e^{-1} (-1)^0 / 1! = 63\%$
  - Trials factor from LEE is  $0.63 \times 10^7$
- How to apply in general?!

# Look-elsewhere effect (Gross&Vitells)



- Approximation best above  $3\sigma$
- Example:
  - $q_{\text{test}} = 4.5$  ( $2.1\sigma$ )
  - 3 crossings at  $0.5\sigma$
  - significance reduced to about  $0.3\sigma$
  - trials factor about 22

Local $\sigma$	Crossings	$\sigma$ ref.	Trials factor	Global $\sigma$
3.5	3	1.0	47	2.3
5.0	3	2.0	290	3.8
7.0	3	2.0	400	6.1

$$q_{\text{test}} = -2 \cdot \text{llr} \approx \Delta\chi^2$$

$$p_0^{\text{global}} \cong p_0^{\text{local}} + \langle N(q_{\text{ref}}) \rangle e^{-(q_{\text{test}} - q_{\text{ref}})/2}$$

# SM Higgs Status @ HCP 18.11.2011


95% CL exclusions:

SINDRUM, Barr et al. ( $\pi, K$  decays):  $\sim 10-110$  MeV

CLEO+CUSB:  $\sim 2 \cdot m_\mu - 5$  GeV



LEP+Tevatron:  $\sim 2 \cdot m_\tau - 114.4$  GeV

Tevatron+ATLAS+CMS: 141-476 GeV




# LHC test of background hypothesis @ HCP

- Dips of  $p_0(\text{local})$ 
  - 119 GeV ( $2.6 \sigma$ )
  - Ignored: 126 GeV ( $2.2 \sigma$ )
  - Ignored: 145 GeV ( $3.1 \sigma$ )
- Roughly (!) 0.9, 0.5, 1.6  $\sigma$  global (up-crossings from data)





ATLAS Combined Higgs Search  
CERN Public Seminar  
December 13, 2011

# SM Higgs production and decay



- +Trigger, bg-rejection, ID-eff.
- Theory uncert. Typically few-20%



Important and fruitful theory-experiment collaboration: [Higgs X-Sec Working Group](#)


# ATLAS Channels in Combination

Higgs decay mode	Mass range (GeV)	Luminosity (1/fb)	
VH->Vbb	-	Not included	
WW->lvqq	240<----->600	1.04 (EPS11) (Back in)	
$\tau\tau$ ->lh3v,ll4v	110<->150	1.06 (LP11)	
ZZ->llqq	200<----->600	2.05	 <b>New</b>
ZZ->llvv	200<----->600	2.05	 <b>New</b>
WW->lvlv	110<----->300	2.05	 <b>New</b>
ZZ->4l	110<----->600	<b>4.8</b>	 <b>New!</b>
$\gamma\gamma$	110<->150	<b>4.9</b>	 <b>New!</b>

# ATLAS Updates




- $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^{(*)} \rightarrow 4l$  quickly analyzed due to relative simplicity
- $H \rightarrow WW \rightarrow lvqq$  back in combination, compliant with LHC Combination Procedures and LHC Cross-Section Working Group Recommendations.
- Other channels will be updated early in 2012.

$$H \rightarrow \gamma\gamma$$





$$H \rightarrow \gamma\gamma$$

- 1.1 -> 4.9/fb
- ~70% of (data-driven) background is SM diphotons
- 9 categories based on  $\gamma \rightarrow e^+e^-$  conversion status, position in the calorimeter,  $p_T^{\gamma\gamma}$  w.r.t  $\gamma\gamma$  thrust axis



# $H \rightarrow \gamma\gamma$ results



- Seen in isolation this channel has, at  $m_H \sim 126$  GeV  
 $p_0$  (local) = 0.27% (2.8  $\sigma$ ) or  
 $p_0$  (global) = 6.5% (1.5  $\sigma$ )
- P.S. LEP hint at 115-116 GeV is threatened




# ATLAS EXPERIMENT

Run Number: 189280,  
Event Number: 143576946  
Date: 2011-09-14, 11:37:11 CET

$$H \rightarrow ZZ^{(*)} \rightarrow 4l$$

EtCut > 0.3 GeV  
PtCut > 3.0 GeV  
Vertex Cuts:  
Z direction < 1cm  
Rphi < 1cm

Muon: blue  
Cells: Tiles, EMC



$$H \rightarrow ZZ^{(*)} \rightarrow 4l$$



- 2.0-2.3  $\rightarrow$  4.8/fb
- Improved e-ID and reco (brem. fitting/GSF)
- For  $m_H=130$  FWHM=4.5 GeV (4 $\mu$ ) – 6.5 GeV (4e)
- For  $m_H=400$  FWHM=35 GeV ( $\sim\Gamma_H$ )
- $m_{4l}$  in likelihood
  - $|m_{12}-m_Z| < 15$  GeV
  - $m_{34} > 15-60$  GeV for  $m_H=120\rightarrow=200$  GeV
- Good mass-resolution and few candidates
  - Structures in  $p_0$  and limits

$$H \rightarrow ZZ^{(*)} \rightarrow 4l$$

	$\mu\mu\mu\mu$		$ee\mu\mu$		$eeee$	
	Low $m_{4\ell}$	High $m_{4\ell}$	Low $m_{4\ell}$	High $m_{4\ell}$	Low $m_{4\ell}$	High $m_{4\ell}$
Int. Luminosity	4.81 fb <sup>-1</sup>		4.81 fb <sup>-1</sup>		4.91 fb <sup>-1</sup>	
$ZZ^{(*)}$	2.0±0.3	16.3±2.4	2.8±0.6	25.2±3.8	1.3±0.3	10.3±1.5
Z, Zbb̄, and t̄t̄	0.16±0.06	0.02±0.01	1.4±0.5	0.17±0.08	1.6±0.7	0.18±0.08
Total Background	2.2±0.3	16.3±2.4	4.2±0.8	25.4±3.8	2.9±0.8	10.5±1.5
Data	3	21	3	27	2	15
$m_H = 125$ GeV	0.58 ± 0.10		0.73 ± 0.13		0.25 ± 0.05	
$m_H = 130$ GeV	1.00 ± 0.17		1.22 ± 0.21		0.43 ± 0.08	
$m_H = 150$ GeV	2.1 ± 0.4		2.9 ± 0.4		1.12 ± 0.18	
$m_H = 200$ GeV	4.9 ± 0.7		7.7 ± 1.0		3.1 ± 0.4	
$m_H = 300$ GeV	2.9 ± 0.4		4.9 ± 0.6		2.1 ± 0.3	
$m_H = 400$ GeV	2.0 ± 0.3		3.3 ± 0.5		1.49 ± 0.21	
$m_H = 600$ GeV	0.34 ± 0.04		0.62 ± 0.10		0.30 ± 0.06	



- “Low mass”:  $100 < m_{4l} < 180$  GeV
- “High mass”:  $180 < m_{4l} < 600$  GeV

$$H \rightarrow ZZ^{(*)} \rightarrow 4l$$





$$H \rightarrow ZZ^{(*)} \rightarrow 4l$$

# signal limits



- No-signal expected exclusion: 137-158, 185 - 400 GeV
- Excluded regions: 135-156, 181-234, 255-415 GeV

# $H \rightarrow ZZ^{(*)} \rightarrow 4l$ background tests



- $p_0$  (local) of 1.8 % (125 GeV), 1.1% (244 GeV), 1.4% (500 GeV)
- $p_0$  (global) > 50% for all peaks if entire mass range considered for LEE, O(30%) for lowest peak if LEE restricted to HCP allowed region (146 GeV @ 99% CL)
- Seen in isolation, none of the excesses of this channel are significant

$$H \rightarrow WW^{(*)} \rightarrow l\nu l\nu$$

- $e^+e^-$ ,  $e^\pm\mu^\mp$ ,  $\mu^+\mu^-$  leptonic final states
- H+0,1 jet exclusive final states (gg, VBF, VH)
- $m_{ll} > 10 \text{ GeV}$  or ( $m_{ll} > 15 \text{ GeV}$  and  $|m_{ll} - m_Z| > 15 \text{ GeV}$ )
- Sliding transverse mass cut
  - $0.75 \cdot m_H < m_T < m_H$  for  $m_H < 220 \text{ GeV}$
  - $0.60 \cdot m_H < m_T < m_H$  for  $m_H > 220 \text{ GeV}$
- Mass-dependent  $\Delta\phi_{ll}$  cut (spin correlations)
  - $\Delta\phi_{ll} < 1.3$  for  $m_H < 170 \text{ GeV}$
  - $\Delta\phi_{ll} < 1.8$  for  $170 \text{ GeV} < m_H < 220 \text{ GeV}$
- $E_{T,rel}^{miss} = E_T^{miss} \cdot \sin(\min(\pi / 2, \Delta\varphi_{miss,l,j})) > 40(ll), 25(ll') \text{ GeV}$





$$H \rightarrow WW^{(*)} \rightarrow l\nu l\nu$$


TABLE I: The expected numbers of signal ( $m_H = 150$  GeV) and background events after the requirements listed in the first column, as well as the observed numbers of events in data. All numbers are summed over lepton flavor.

$H + 0$ -jet Channel	Signal	$WW$	$W + \text{jets}$	$Z/\gamma^* + \text{jets}$	$t\bar{t}$	$tW/tb/tqb$	$WZ/ZZ/W\gamma$	Total Bkg.	Observed
Jet Veto	$99 \pm 21$	$524 \pm 52$	$84 \pm 41$	$174 \pm 169$	$42 \pm 14$	$32 \pm 8$	$15 \pm 4$	$872 \pm 182$	920
$p_T^{\ell\ell} > 30$ GeV	$95 \pm 20$	$467 \pm 45$	$69 \pm 34$	$30 \pm 12$	$39 \pm 14$	$29 \pm 8$	$13 \pm 4$	$648 \pm 60$	700
$m_{\ell\ell} < 50$ GeV	$68 \pm 15$	$118 \pm 15$	$21 \pm 8$	$13 \pm 8$	$7 \pm 4$	$5.8 \pm 1.8$	$1.9 \pm 0.6$	$166 \pm 19$	199
$\Delta\phi_{\ell\ell} < 1.3$	$58 \pm 13$	$91 \pm 12$	$12 \pm 5$	$9 \pm 6$	$6 \pm 3$	$5.8 \pm 1.8$	$1.7 \pm 0.6$	$125 \pm 15$	149
$0.75 m_H < m_T < m_H$	$40 \pm 9$	$52 \pm 7$	$5 \pm 2$	$2 \pm 4$	$2.4 \pm 1.6$	$1.5 \pm 1.0$	$1.1 \pm 0.5$	$63 \pm 9$	81
$H + 1$ -jet Channel	Signal	$WW$	$W + \text{jets}$	$Z/\gamma^* + \text{jets}$	$t\bar{t}$	$tW/tb/tqb$	$WZ/ZZ/W\gamma$	Total Bkg.	Observed
1 jet	$50 \pm 9$	$193 \pm 20$	$38 \pm 21$	$74 \pm 65$	$473 \pm 124$	$174 \pm 26$	$14 \pm 2$	$967 \pm 145$	952
$b$ -jet veto	$48 \pm 9$	$188 \pm 19$	$35 \pm 19$	$73 \pm 61$	$174 \pm 49$	$66 \pm 11$	$14 \pm 2$	$549 \pm 83$	564
$ \mathbf{p}_T^{\text{tot}}  < 30$ GeV	$39 \pm 7$	$154 \pm 16$	$18 \pm 9$	$38 \pm 32$	$106 \pm 30$	$50 \pm 9$	$9.7 \pm 1.5$	$376 \pm 48$	405
$Z \rightarrow \tau\tau$ veto	$39 \pm 7$	$150 \pm 17$	$18 \pm 8$	$34 \pm 23$	$102 \pm 23$	$48 \pm 8$	$9 \pm 2$	$361 \pm 38$	388
$m_{\ell\ell} < 50$ GeV	$26 \pm 6$	$33 \pm 5$	$3.3 \pm 1.4$	$8 \pm 7$	$20 \pm 7$	$11 \pm 3$	$1.8 \pm 0.5$	$77 \pm 12$	90
$\Delta\phi_{\ell\ell} < 1.3$	$23 \pm 5$	$25 \pm 4$	$2.1 \pm 1.0$	$4 \pm 6$	$17 \pm 6$	$9 \pm 3$	$1.5 \pm 0.4$	$60 \pm 10$	72
$0.75 m_H < m_T < m_H$	$14 \pm 3$	$12 \pm 3$	$0.9 \pm 0.4$	$1.3 \pm 1.9$	$8 \pm 2$	$4.0 \pm 1.6$	$0.7 \pm 0.3$	$28 \pm 4$	29
Control Regions	Signal	$WW$	$W + \text{jets}$	$Z/\gamma^* + \text{jets}$	$t\bar{t}$	$tW/tb/tqb$	$WZ/ZZ/W\gamma$	Total Bkg.	Observed
$WW$ 0-jet ( $m_H < 220$ GeV)	$1.7 \pm 0.4$	$223 \pm 30$	$20 \pm 15$	$6 \pm 8$	$25 \pm 10$	$15 \pm 4$	$8 \pm 3$	$296 \pm 36$	296
$WW$ 0-jet ( $m_H \geq 220$ GeV)	$10 \pm 2$	$173 \pm 23$	$24 \pm 12$	$13 \pm 19$	$15 \pm 6$	$8 \pm 3$	$3.3 \pm 0.6$	$236 \pm 33$	258
$WW$ 1-jet ( $m_H < 220$ GeV)	$1.0 \pm 0.3$	$76 \pm 13$	$5 \pm 3$	$5 \pm 5$	$56 \pm 14$	$23 \pm 5$	$5.3 \pm 1.4$	$171 \pm 21$	184
$WW$ 1-jet ( $m_H \geq 220$ GeV)	$5.8 \pm 1.5$	$51 \pm 9$	$3.9 \pm 1.8$	$10 \pm 10$	$35 \pm 9$	$18 \pm 4$	$2.8 \pm 0.6$	$120 \pm 17$	129
$t\bar{t}$ 1-jet	$0.9 \pm 0.3$	$3.9 \pm 1.0$	-	$1 \pm 17$	$184 \pm 64$	$80 \pm 19$	$0.2 \pm 0.9$	$270 \pm 69$	249

$$H \rightarrow WW^{(*)} \rightarrow l\nu l\nu$$





- Largest excess  $1.9 \sigma$
- No-signal expected 95% CL exclusion in range 134-200 GeV
- Exclusion at  $\geq 95\%$  CL in region 145-206 GeV



$$H \rightarrow ZZ \rightarrow l^+ l^- \nu \bar{\nu}, l = e, \mu$$

- Some similarity to  $H \rightarrow WW^{(*)} \rightarrow l\nu l\nu$   
but no  $e\mu$  channel and  $|m_Z - m_W| < 15 \text{ GeV}$

$$H \rightarrow ZZ \rightarrow l^+ l^- \nu_l \bar{\nu}_l, l = e, \mu$$






	$m_H < 280$ GeV	$m_H > 280$ GeV
ETmiss	$> 66$ GeV	$> 82$ GeV
Lepton opening angle	$1 < \Delta\phi_{  } < 2.64$	$\Delta\phi_{  } < 2.25$

$$H \rightarrow ZZ \rightarrow l^+ l^- \nu_l \bar{\nu}_l, l = e, \mu$$

Source	low $m_H$ search	high $m_H$ search	
Z + jets	$40.5 \pm 4.1 \pm 3.1$	$9.9 \pm 2.1 \pm 3.6$	
W + jets	$17 \pm 4 \pm 17$	$6.1 \pm 1.7 \pm 6.1$	
top	$40 \pm 1 \pm 12$	$20.9 \pm 1.0 \pm 6.2$	
Multijet	$1.2 \pm 0.7 \pm 0.6$	$0.4 \pm 0.4 \pm 0.2$	
ZZ	$35.1 \pm 0.7 \pm 4.1$	$29.6 \pm 0.6 \pm 3.5$	
WZ	$32.4 \pm 1.0 \pm 3.8$	$23.2 \pm 0.8 \pm 2.7$	
WW	$25.4 \pm 0.7 \pm 3.1$	$9.4 \pm 0.4 \pm 1.1$	
Total BG	$192 \pm 6 \pm 35$	$100 \pm 3 \pm 17$	
Data	175	89	
$m_H$ [GeV]	Signal expectation		$s/b$
200	$9.9 \pm 0.2 \pm 1.8$		8%
300	$19.9 \pm 0.3 \pm 3.5$		24%
400	$19.6 \pm 0.3 \pm 3.4$		57%
500	$8.8 \pm 0.1 \pm 1.5$		60%
600	$3.6 \pm 0.0 \pm 0.6$		60%


$$H \rightarrow ZZ \rightarrow l^+ l^- \nu_l \bar{\nu}_l, l = e, \mu$$







- Fit to  $m_T$  distribution
- No-signal expected 95% CL exclusion between 320-460 GeV
- 95% CL excluded region 310-470 GeV
- Impact on limit of uncertainty in off-shell Higgs production  $\sim 3$ -20% for 200-600 GeV

$$H \rightarrow ZZ \rightarrow l^+ l^- q \bar{q}$$

- Small missing  $E_T$
- Leptonic+hadronic Z decays
- Z+jets background, Z→bb  
20%: 2b-tagged and untagged categories
  - $\epsilon_b \sim 70\%$ , light q-rejection  $\sim 100$
- Z's boosted for  $m_H > 300$  GeV  
("low/high mass" searches)









$$H \rightarrow ZZ \rightarrow l^+ l^- q \bar{q}$$



$$H \rightarrow ZZ \rightarrow l^+ l^- q \bar{q}$$


b-Tagged



Low-mass




High-mass

$$H \rightarrow ZZ \rightarrow l^+ l^- q \bar{q}$$




- $p_0 = 0.13 - 0.5$  (no sig. excesses)
- Max signal sensitivity @ 360 GeV
  - 1.7xSM expected limit
  - 1.2xSM excluded @ 95% CL
- Limits 1.2-12xSM 200-600 GeV

# $H \rightarrow WW \rightarrow l\nu jj$



- No excesses
- Maximum signal sensitivity  
4.5 x SM @ 400 GeV
- Exclude 2.7-20 x SM  
between 250-600 GeV

$$H \rightarrow \tau^+ \tau^- \rightarrow l^+ l^- 4\nu$$




	$ee + \mu\mu + e\mu$
$Z/\gamma^* \rightarrow \tau^+ \tau^-$	$25.4 \pm 2.7$
$Z/\gamma^* \rightarrow \ell^+ \ell^-$ ( $\ell=e,\mu$ )	$3.7 \pm 1.2$
$t\bar{t}$	$13.2 \pm 2.2$
Single- $t$	$1.2 \pm 0.5$
Di-boson	$1.6 \pm 0.6$
Backgrounds with fake Leptons	$2.2 \pm 0.9$
Total Background expectation	$47.4 \pm 3.9$
Observed data	46



  

	SM signal	$gg \rightarrow H$	VBF
$m_H = 110$ GeV	$0.39 \pm 0.06$	$0.35 \pm 0.02$	
$m_H = 115$ GeV	$0.39 \pm 0.06$	$0.35 \pm 0.02$	
$m_H = 120$ GeV	$0.44 \pm 0.05$	$0.38 \pm 0.02$	
$m_H = 130$ GeV	$0.40 \pm 0.04$	$0.33 \pm 0.01$	
$m_H = 140$ GeV	$0.21 \pm 0.02$	$0.19 \pm 0.01$	


- 2 leptons, 1 jet (VBF, gg-NLO), missing energy
- H-boost enhances S/B after topological selection
- Collinear approximation used to estimate  $m_{\tau\tau}$  ( $\nu$ 's // to  $l$ 's)
- Count events in mass window 100-150 GeV
- Irreducible Z- $\tau\tau$  estimated by embedding simulated  $\tau$ 's into  $\mu^+\mu^-$ (Z) events
- Exclusion limits  $\sim 30\times$ SM obtained



$$H \rightarrow \tau^+ \tau^- \rightarrow lh3\nu$$




- 1 leptonic, 1 hadronic decay, missing  $E_T$
- MMC (Missing Mass Calculator), allows non-collinear  $\nu$ 's, mode of  $m_{\tau\tau}$  from scanning over under-determined kinematic variables, weighted by pdf of 3D opening angle between visible and invisible decay products
- Exclusion 7-26 x SM (expected 12-30)



# Combined Results

$$L(m_H, \mu, \vec{\vartheta}) = \prod_i L_i(m_H, \mu, \vec{\vartheta}_i)$$



# Correlated syst. uncert. on signal yield (%)

for a Higgs boson mass hypothesis of 120 GeV for the  $H \rightarrow \gamma\gamma$ ,  $H \rightarrow \tau\tau \rightarrow \ell\tau_{had}3\nu$ ,  $H \rightarrow \tau\tau \rightarrow \ell^+\ell^-4\nu$ ,  $H \rightarrow WW^{(*)} \rightarrow \ell^+\nu\ell^-\bar{\nu}$  and  $H \rightarrow ZZ^{(*)} \rightarrow \ell^+\ell^-\ell^+\ell^-$  analyses and 300 GeV for the  $H \rightarrow ZZ \rightarrow \ell^+\ell^-q\bar{q}$ ,  $H \rightarrow ZZ \rightarrow \ell^+\ell^-\nu\bar{\nu}$  and  $H \rightarrow WW \rightarrow \ell\nu q\bar{q}'$  analyses.

	$H \rightarrow \tau^+\tau^-$		$H \rightarrow \gamma\gamma$	$H \rightarrow WW^{(*)}$		$H \rightarrow ZZ^{(*)}$		
	$\ell\tau_{had}3\nu$	$\tau_\ell\tau_\ell + jet$		$\ell\nu\ell\nu$	$\ell\nu q\bar{q}$	$llll$	$ll\nu\nu$	$llq\bar{q}$
Luminosity	+3.8 -3.6	+3.8 -3.6	+4.0 -3.8	+3.8 -3.6	+3.8 -3.6	+3.9 -3.8	+3.8 -3.6	+3.8 -3.6
$e/\gamma$ eff.	$\pm 3.5$	$\pm 2.0$	$^{+13.5}_{-11.9}$	$\pm 2.0$	$\pm 0.9$	$\pm 2.9$	$\pm 1.2$	$\pm 1.2$
$e/\gamma$ E. scale	$^{+1.3}_{-0.1}$	$\pm 0.3$	-	$\pm 0.4$	-	-	$\pm 0.7$	$\pm 0.4$
$e/\gamma$ res.	-	$^{+0.2}_{-0.5}$	-	$^{+0.20}_{-0.05}$	-	-	$\pm 0.25$	$\pm 0.1$
$\mu$ eff.	$\pm 1.0$	$\pm 2.0$	-	-	$\pm 0.3$	$\pm 0.16$	$\pm 0.7$	$\pm 0.5$
$\mu$ res. Id.	-	$^{+0.2}_{-0.5}$	-	$^{+0.02}_{-0.04}$	-	-	$\pm 1.1$	$\pm 1.1$
$\mu$ res. MS.	-	-	-	$^{+0.04}_{+0.08}$	-	-	$^{+1.1}_{-1.0}$	$\pm 1.1$
Jet/ $\tau$ /MET E. scale	+18.9 -16.4	+3.4 -10.0	-	+4.46 -6.47	+18.4 -15.5	-	$\pm 1.6$	$\pm 15.0$
JER	-	$\pm 2.0$	-	+1.8 -1.7	+9.0 -8.2	-	+0.3 -0.0	+4.0 -0.0
MET	-	+4.4 -5.3	-	+1.8 -1.7	-	-	-	-
$b$ -tag eff.	-	-	-	$\pm 0.5$	-	-	$\pm 0.3$	$\pm 3.7$
$\tau$ eff.	$\pm 9.1$	-	-	-	-	-	-	-

Note: Signal uncertainties affect limit, less important for testing background.


# Correlated syst. uncert. (relative) on background

	$H \rightarrow \tau^+ \tau^-$		$H \rightarrow \gamma\gamma$	$H \rightarrow WW^{(*)}$		$H \rightarrow ZZ^{(*)}$		
	$\ell\tau_{had}3\nu$	$\tau_\ell\tau_\ell + jet$		$\ell\nu\ell\nu$	$\ell\nu qq$	$llll$	$ll\nu\nu$	$llqq$
Luminosity	+3.0 -2.9	+3.8 -3.6	-	$\pm 0.2$	-	+3.7 -3.6	+2.4 -2.3	+0.3 -0.2
$e/\gamma$ eff.	$\pm 2.4$	+0.5 -1.6	-	$\pm 2.3$	$\pm 0.8$	$\pm 1.6$	$\pm 0.8$	$\pm 0.1$
$e/\gamma$ E. scale	+0.9 -0.3	$\pm 0.8$	-	+0.2 -0.1	-	-	+1.7 -1.6	$\pm 0.1$
$e/\gamma$ res.	-	+0.3 -2.6	-	+0.1 -0.0	-	-	$\pm 0.6$	$\pm 0.2$
$\mu$ eff.	$\pm 1.4$	+0.5 -1.6	-	-	$\pm 0.3$	$\pm 0.1$	$\pm 0.5$	$\pm 0.03$
$\mu$ res. Id.	-	+0.3 -2.6	-	-0.03 -0.06	-	-	+1.7 -1.6	$\pm 0.2$
$\mu$ res. MS.	-	-	-	+0.00 -0.02	-	-	+1.7 -1.6	$\pm 0.2$
Jet/ $\tau$ /MET E. scale	+10.0 -8.9	+7.0 -9.8	-	+8.5 -10.4	-	-	+6.9 -5.2	$\pm 1.0$
JER	-	$\pm 2.5$	-	+3.3 -3.0	-	-	+1.8 -0.0	+0.3 -0.0
MET	-	+0.4 -2.7	-	+0.6 -0.5	-	-	-	-
$b$ -tag eff.	-	-	-	$\pm 1.8$	-	-	+7.0 -5.5	$\pm 0.2$
$\tau$ eff.	$\pm 7.2$	-	-	-	-	-	-	-



for a Higgs boson mass hypothesis of 120 GeV for the  $H \rightarrow \gamma\gamma$ ,  $H \rightarrow \tau\tau \rightarrow \ell\tau_{had}3\nu$ ,  $H \rightarrow \tau\tau \rightarrow \ell^+\ell^-4\nu$ ,  $H \rightarrow WW^{(*)} \rightarrow \ell^+\nu\ell^-\bar{\nu}$  and  $H \rightarrow ZZ^{(*)} \rightarrow \ell^+\ell^-\ell^+\ell^-$  analyses and 300 GeV for the  $H \rightarrow ZZ \rightarrow \ell^+\ell^-q\bar{q}$ ,  $H \rightarrow ZZ \rightarrow \ell^+\ell^-\nu\bar{\nu}$  and  $H \rightarrow WW \rightarrow \ell\nu q\bar{q}'$  analyses.

Note: Important for testing background hypothesis.

# CLs limits all channels



# CLs – test of SM signal hypothesis





Expected exclusion region: 125-520 GeV



Observed exclusions: 112.7-115.5, 131-237, 251-453 GeV

(LHC@HCP: Expected 124-520, Observed 141-476)

# CLs upper limits on $\sigma/\sigma_{\text{SM}}$





# $p_0$ – test of background hypothesis



- At 126 GeV local  $p_0=1.9 \times 10^{-4}$  ( $3.6 \sigma$ )
- 6 crossings in full mass range (see also Cyan Band plots)
  - Global  $p_0=1.1\%$  ( $2.3 \sigma$ )





# Estimate of signal strength $\mu = \sigma/\sigma_{SM}$




- Allowing negative values of  $\mu$  using a specific treatment to not allow pdfs to have negative values.
- Don't use these to set limits or test background hypothesis, they are for illustration.

Signal strength  $\sim 1.5$  around 126 GeV is not incompatible with SM Higgs expectation of 1

# 3 most sensitive channels




# Asymptotics (baseline) vs. MC ensemble



# Next steps

- Finalize 2011 results
- Combine published ATLAS+CMS 2011 results
  - Inter-experiment energy scale systematics important
- Prepare improved analyses for 2012
  - $\sim 4$  times data and a bit higher  $\sqrt{s}$  could take ATLAS+CMS within spitting distance of  $5\sigma$  or exclude present excess as fluctuation
  - Should decide on LEE-region before the data arrive
- If  $\sim 124\text{-}6$  GeV persists:
  - Measurement of mass (ML, not  $M\sigma$ )
  - $\gamma\gamma$ : It's not a vector
  - $\tau\tau$ ,  $bb$ ,  $WW$ ,  $ZZ$  decays all accessible for measurements
  - Consequences for BSM...
- If not:
  - Go down in the couplings
  - Don't forget (difficult) high mass



# Conclusions

- Preliminary results for SM Higgs boson searches with ATLAS in up to 4.9/fb of 2011 data
- Expected exclusion region  $m_H=124.6-520$  GeV
- 3 regions excluded at  $\geq 95\%$  CL
  - 112.7-115.5, 131-237, 251-453 GeV
- Excesses observed in 3 most sensitive channels near  $m_H=126$  GeV (local: 2.7, 2.0, 1.4  $\sigma$ )
  - Global probability for background to give this combined excess is approximately 1%