

about R-Hadrons & how to find them

A (nearly) published analysis.

Simon Heisterkamp

May 2012

What are R-hadrons

- collective name for heavy stable long lived particles hadronized with SM particles


- e.g. gluinos, squarks,

- many SUSY models predict them

- long lifetime!

- results from small mass difference to LSP


- necessary for hadronization



How are they observed

- Long lifetime \rightarrow direct observation
- Very high mass (TeV scale) \rightarrow slow ($\beta < 1$)
 - time of flight measurements
 - specific ionization (dE/dx) relates to $\beta\gamma$ via Bethe-Bloch equation.

How are they observed



How are they observed

- ToF measured by:
 - calorimeters
 - muon spectrometer
- dE/dx measured by:
 - Pixel detector (ADC readout)
 - TRT (Time over threshold, high threshold)
 - Calorimeters

Phenomenology



- various quark systems possible:
 - R-mesons & -baryons
 - charges $-2, -1, 0, 1, 2$
- Interaction model:
 - Stable & massive: Direct cross-section is tiny.
 - Associated light quark system interacts like a hadron with a big momentum reservoir

Independent of SUSY
model

Performance


Pixel performance

-calibrate with protons





Performance

Calorimeter performance





MC agreement
is terrible
→ calibration



Performance

MS performance



Limits

