Contribution ID: 2 Type: not specified

Reconstruction of meriodional flow speed of a solar dynamo model by data assimilation algorithm

Monday, 9 November 2015 15:00 (30 minutes)

Accurate knowledge of time-variation in meridional flow-speed and profile is crucial for estimating a solar cycle's features, which are ultimately responsible for causing space climate variations. However, no consensus has been reached yet about the Sun's meridional circulation pattern observations and theories. By implementing an Ensemble Kalman Filter (EnKF) data assimilation in a Babcock-Leighton solar dynamo model using Data Assimilation Research Testbed (DART) framework, we find that the best reconstruction of time-variation in meridional flow-speed can be obtained when ten or more observations are used with an updating time of 15 days and a $\leq 10\%$ observational error. Increasing ensemble-size from 16 to 160 improves reconstruction. Comparison of reconstructed flow-speed with "true-state" reveals that EnKF data assimilation is very powerful for reconstructing meridional flow-speeds and suggests that it can be implemented for reconstructing spatio-temporal patterns of meridional circulation.

Primary author: Dr MITRA, Dhrubaditya (NORDITA (Nordic Institute of Theoretical Physics), Stockholm)

Presenter: Dr MITRA, Dhrubaditya (NORDITA (Nordic Institute of Theoretical Physics), Stockholm)