


Exclusive Radiative Higgs Decays in the Standard Model and Beyond

Stefan Alte, Johannes Gutenberg-Universität Mainz

Matthias König and Matthias Neubert



HEFT2016, Copenhagen

26 October 2016





Introduction

Back in the year 2012 ...

Introduction



Back in the year 2012 ...



... the **final piece in the SM puzzle** - the **Higgs boson** - was discovered

Introduction

Back in the year 2012 ...



... the **final piece in the SM puzzle** - the **Higgs boson** - was discovered

The path to new physics may open up in the **properties of the Higgs boson** due to its **key role in the SM!**

Introduction

The **dominant decay channels** are studied with **large effort**

Introduction

The **dominant decay channels** are studied with **large effort**

Channels which are **subdominant or absent in the SM** could be **dominated by NP effects!**

Introduction

The **dominant decay channels** are studied with **large effort**

Channels which are **subdominant or absent in the SM** could be **dominated by NP effects!**

We focus on the **exclusive radiative decays** $h \rightarrow MV$ into a **meson M** and a gauge boson $V = \{\gamma, Z, W\}$

Introduction

The **dominant decay channels** are studied with **large effort**

Channels which are **subdominant or absent in the SM** could be **dominated by NP effects!**

We focus on the **exclusive radiative decays** $h \rightarrow MV$ into a **meson M** and a gauge boson $V = \{\gamma, Z, W\}$

“Exclusive Radiative Higgs Decays as Probes of Light-Quark Yukawa Couplings”, König and Neubert (2015), JHEP **1508** (2015) 012, arXiv: **1505.03870**

“Exclusive Weak Radiative Higgs Decays in the Standard Model and Beyond”, Alte, König and Neubert (2016), arXiv: **1609.06310**

Effective Higgs Couplings

$$\begin{aligned}
 \mathcal{L}_{\text{eff}}^{\text{Higgs}} = & \kappa_W \frac{2m_W^2}{v} h W_\mu^+ W^{-\mu} + \kappa_Z \frac{m_Z^2}{v} h Z_\mu Z^\mu - \frac{h}{\sqrt{2}} \sum_{f=u,d,e} (\bar{f}_L Y_f f_R + \text{h.c.}) \\
 & + \frac{\alpha}{4\pi v} \left(\kappa_{\gamma\gamma} h F_{\mu\nu} F^{\mu\nu} - \tilde{\kappa}_{\gamma\gamma} h F_{\mu\nu} \tilde{F}^{\mu\nu} \right. \\
 & \left. + \frac{2\kappa_{\gamma Z}}{\sin \theta_W \cos \theta_W} h F_{\mu\nu} Z^{\mu\nu} - \frac{2\tilde{\kappa}_{\gamma Z}}{\sin \theta_W \cos \theta_W} h F_{\mu\nu} \tilde{Z}^{\mu\nu} \right)
 \end{aligned}$$

$(Y_f)_{ii} = (\kappa_{f_i} + i\tilde{\kappa}_{f_i}) \frac{\sqrt{2}m_{f_i}}{v}$

Effective Higgs Couplings

$$\begin{aligned}\mathcal{L}_{\text{eff}}^{\text{Higgs}} = & \kappa_W \frac{2m_W^2}{v} h W_\mu^+ W^{-\mu} + \kappa_Z \frac{m_Z^2}{v} h Z_\mu Z^\mu - \frac{h}{\sqrt{2}} \sum_{f=u,d,e} (\bar{f}_L Y_f f_R + \text{h.c.}) \\ & + \frac{\alpha}{4\pi v} \left(\kappa_{\gamma\gamma} h F_{\mu\nu} F^{\mu\nu} - \tilde{\kappa}_{\gamma\gamma} h F_{\mu\nu} \tilde{F}^{\mu\nu} \right. \\ & \left. + \frac{2\kappa_{\gamma Z}}{\sin\theta_W \cos\theta_W} h F_{\mu\nu} Z^{\mu\nu} - \frac{2\tilde{\kappa}_{\gamma Z}}{\sin\theta_W \cos\theta_W} h F_{\mu\nu} \tilde{Z}^{\mu\nu} \right)\end{aligned}$$

$(Y_f)_{ii} = (\kappa_{f_i} + i\tilde{\kappa}_{f_i}) \frac{\sqrt{2}m_{f_i}}{v}$

SM: blue parameters $\rightarrow 1$, red parameters $\rightarrow 0$, $Y_f \rightarrow$ diagonal,
 $\kappa_{f_i} \rightarrow 1$, $\tilde{\kappa}_{f_i} \rightarrow 0$

Effective Higgs Couplings

$$\begin{aligned}\mathcal{L}_{\text{eff}}^{\text{Higgs}} = & \kappa_W \frac{2m_W^2}{v} h W_\mu^+ W^{-\mu} + \kappa_Z \frac{m_Z^2}{v} h Z_\mu Z^\mu - \frac{h}{\sqrt{2}} \sum_{f=u,d,e} (\bar{f}_L Y_f f_R + \text{h.c.}) \\ & + \frac{\alpha}{4\pi v} \left(\kappa_{\gamma\gamma} h F_{\mu\nu} F^{\mu\nu} - \tilde{\kappa}_{\gamma\gamma} h F_{\mu\nu} \tilde{F}^{\mu\nu} \right. \\ & \left. + \frac{2\kappa_{\gamma Z}}{\sin\theta_W \cos\theta_W} h F_{\mu\nu} Z^{\mu\nu} - \frac{2\tilde{\kappa}_{\gamma Z}}{\sin\theta_W \cos\theta_W} h F_{\mu\nu} \tilde{Z}^{\mu\nu} \right)\end{aligned}$$

$(Y_f)_{ii} = (\kappa_{f_i} + i\tilde{\kappa}_{f_i}) \frac{\sqrt{2}m_{f_i}}{v}$

SM: blue parameters $\rightarrow 1$, red parameters $\rightarrow 0$, $Y_f \rightarrow$ diagonal,
 $\kappa_{f_i} \rightarrow 1$, $\tilde{\kappa}_{f_i} \rightarrow 0$

The Lagrangian $\mathcal{L}_{\text{eff}}^{\text{Higgs}}$ allows for a **model-independent analysis** of the Higgs couplings

Exclusive Radiative Decays $h \rightarrow M\gamma$

König and Neubert (2015)

Light-Quark Yukawa Couplings

Idea: access the **light-quark Yukawa couplings** ($q \neq t$) in the decays $h \rightarrow M\gamma$ Bodwin et al. (2013); Kagan et al. (2015)

Light-Quark Yukawa Couplings

Idea: access the **light-quark Yukawa couplings** ($q \neq t$) in the decays $h \rightarrow M\gamma$ Bodwin et al. (2013); Kagan et al. (2015)


Current LHC measurements: no strong constraints on the parameters κ_q Kagan et al. (2015); Perez et al. (2015)

$$\sqrt{|\kappa_u|^2 + |\tilde{\kappa}_u|^2} < 3000$$


$$\sqrt{|\kappa_d|^2 + |\tilde{\kappa}_d|^2} < 1500$$

$$\sqrt{|\kappa_s|^2 + |\tilde{\kappa}_s|^2} < 75$$

Decay Topologies




Decay Topologies



The **direct contribution** involves κ_q and $\tilde{\kappa}_q$ and is calculated within the **QCD factorisation approach** Brodsky, Lepage (1979); Efremov, Radyushkin (1980); Chernyak, Zhitnitsky (1984)


Decay Topologies



The **direct contribution** involves κ_q and $\tilde{\kappa}_q$ and is calculated within the **QCD factorisation approach** Brodsky, Lepage (1979); Efremov, Radyushkin (1980); Chernyak, Zhitnitsky (1984)

The **indirect contribution** involves an **off-shell gauge boson** which **converts into the meson**


Decay Topologies



The **direct contribution** involves κ_q and $\tilde{\kappa}_q$ and is calculated within the **QCD factorisation approach** Brodsky, Lepage (1979); Efremov, Radyushkin (1980); Chernyak, Zhitnitsky (1984)

The **indirect contribution** involves an **off-shell gauge boson** which **converts into the meson**

The **effective vertex** includes the **SM contributions** Bergström and Hulth (1985)



as well as the **NP contributions** $\kappa_{\gamma\gamma}$ and $\kappa_{\gamma Z}$

Decay Amplitude

$$i\mathcal{A}(h \rightarrow M\gamma) = -\frac{ef_M}{2} \left[\left(\varepsilon_M^* \cdot \varepsilon_\gamma^* - \frac{q \cdot \varepsilon_M^* k \cdot \varepsilon_\gamma^*}{k \cdot q} \right) F_1^M - i\epsilon_{\mu\nu\alpha\beta} \frac{k^\mu q^\nu \varepsilon_M^{*\alpha} \varepsilon_\gamma^{*\beta}}{k \cdot q} F_2^M \right]$$

Decay Amplitude

$$i\mathcal{A}(h \rightarrow M\gamma) = -\frac{ef_M}{2} \left[\left(\epsilon_M^* \cdot \epsilon_\gamma^* - \frac{q \cdot \epsilon_M^* k \cdot \epsilon_\gamma^*}{k \cdot q} \right) F_1^M - i\epsilon_{\mu\nu\alpha\beta} \frac{k^\mu q^\nu \epsilon_M^{*\alpha} \epsilon_\gamma^{*\beta}}{k \cdot q} F_2^M \right]$$

The **form factors** receive **direct and indirect** contributions

Decay Amplitude

$$i\mathcal{A}(h \rightarrow M\gamma) = -\frac{ef_M}{2} \left[\left(\varepsilon_M^* \cdot \varepsilon_\gamma^* - \frac{q \cdot \varepsilon_M^* k \cdot \varepsilon_\gamma^*}{k \cdot q} \right) F_1^M - i\epsilon_{\mu\nu\alpha\beta} \frac{k^\mu q^\nu \varepsilon_M^{*\alpha} \varepsilon_\gamma^{*\beta}}{k \cdot q} F_2^M \right]$$

The **form factors** receive **direct and indirect** contributions

Neglecting numerically small terms, one finds for a J/ψ meson

$$F_1^{J/\psi} = -0.137\kappa_W + 0.030\kappa_t + 0.033\kappa_{\gamma\gamma} + 0.007\kappa_c + \dots$$

Decay Amplitude

$$i\mathcal{A}(h \rightarrow M\gamma) = -\frac{ef_M}{2} \left[\left(\varepsilon_M^* \cdot \varepsilon_\gamma^* - \frac{q \cdot \varepsilon_M^* k \cdot \varepsilon_\gamma^*}{k \cdot q} \right) F_1^M - i\epsilon_{\mu\nu\alpha\beta} \frac{k^\mu q^\nu \varepsilon_M^{*\alpha} \varepsilon_\gamma^{*\beta}}{k \cdot q} F_2^M \right]$$

The **form factors** receive **direct and indirect** contributions

Neglecting numerically small terms, one finds for a J/ψ meson

$$F_1^{J/\psi} = -0.137\kappa_W + 0.030\kappa_t + 0.033\kappa_{\gamma\gamma} + 0.007\kappa_c + \dots$$

It is challenging to access κ_c since the dominant terms arise from the indirect contribution

Reduction of Theoretical Uncertainties

Need to **know** the indirect contribution **very precisely**

Reduction of Theoretical Uncertainties

Need to **know** the indirect contribution **very precisely**

Subtract the indirect contribution **without assuming that SM is correct**

Reduction of Theoretical Uncertainties

Need to **know** the indirect contribution **very precisely**

Subtract the indirect contribution **without assuming that SM is correct**



Reduction of Theoretical Uncertainties

Need to **know** the indirect contribution **very precisely**

Subtract the indirect contribution **without assuming that SM is correct**



Normalise $\Gamma(h \rightarrow M\gamma)$ to $\Gamma(h \rightarrow \gamma\gamma)$: $\frac{\Gamma(h \rightarrow M\gamma)}{\Gamma(h \rightarrow \gamma\gamma)} = \frac{\text{Br}(h \rightarrow M\gamma)}{\text{Br}(h \rightarrow \gamma\gamma)}$

Reduction of Theoretical Uncertainties

Need to **know** the indirect contribution **very precisely**

Subtract the indirect contribution **without assuming that SM is correct**





Normalise $\Gamma(h \rightarrow M\gamma)$ to $\Gamma(h \rightarrow \gamma\gamma)$: $\frac{\Gamma(h \rightarrow M\gamma)}{\Gamma(h \rightarrow \gamma\gamma)} = \frac{\text{Br}(h \rightarrow M\gamma)}{\text{Br}(h \rightarrow \gamma\gamma)}$

Additional benefit: no dependence on Γ_h



Branching Ratios as Functions of κ_q

Assumption: no CP-odd coupling ($\tilde{\kappa}_q \rightarrow 0$)



Branching Ratios as Functions of κ_q



Assumption: no CP-odd coupling ($\tilde{\kappa}_q \rightarrow 0$)



$\text{Br}(h \rightarrow J/\psi\gamma)$ exhibits a **mild dependence on κ_c**

Branching Ratios as Functions of κ_q

Assumption: no CP-odd coupling ($\tilde{\kappa}_q \rightarrow 0$)





$\text{Br}(h \rightarrow J/\psi\gamma)$ exhibits a **mild dependence** on κ_c

$\text{Br}(h \rightarrow \Upsilon(1S)\gamma)$ depends **strongly** on κ_b due to a **fortuitous cancellation** of the direct and the indirect contribution



$\text{Br}(h \rightarrow \Upsilon\gamma)$ as Functions of κ_b and $\tilde{\kappa}_b$

Possible future scenarios



$\text{Br}(h \rightarrow \Upsilon\gamma)$ as Functions of κ_b and $\tilde{\kappa}_b$

Possible future scenarios





Blue: measurements of $h \rightarrow b\bar{b}$

Orange: measurements of $h \rightarrow \Upsilon\gamma$

Br($h \rightarrow \Upsilon\gamma$) as Functions of κ_b and $\tilde{\kappa}_b$

Possible future scenarios



Blue: measurements of $h \rightarrow b\bar{b}$

Orange: measurements of $h \rightarrow \Upsilon\gamma$

The **overlap** can provide **information** about the **CP nature**

Branching Ratios in the SM

$$\text{BR}(h \rightarrow \rho^0 \gamma) = (1.68 \pm 0.02_f \pm 0.08_{h \rightarrow \gamma\gamma}) \cdot 10^{-5}$$

$$\text{BR}(h \rightarrow \omega \gamma) = (1.48 \pm 0.03_f \pm 0.07_{h \rightarrow \gamma\gamma}) \cdot 10^{-6}$$

$$\text{BR}(h \rightarrow \phi \gamma) = (2.31 \pm 0.03_f \pm 0.11_{h \rightarrow \gamma\gamma}) \cdot 10^{-6}$$

$$\text{BR}(h \rightarrow J/\psi \gamma) = (2.95 \pm 0.07_f \pm 0.06_{\text{direct}} \pm 0.14_{h \rightarrow \gamma\gamma}) \cdot 10^{-6}$$

$$\text{BR}(h \rightarrow \Upsilon(1S) \gamma) = (4.61 \pm 0.06_{f_{-1.21}^{+1.75}} \pm 0.22_{\text{direct}} \pm 0.22_{h \rightarrow \gamma\gamma}) \cdot 10^{-9}$$

$$\text{BR}(h \rightarrow \Upsilon(2S) \gamma) = (2.34 \pm 0.04_{f_{-0.99}^{+0.75}} \pm 0.11_{\text{direct}} \pm 0.11_{h \rightarrow \gamma\gamma}) \cdot 10^{-9}$$

$$\text{BR}(h \rightarrow \Upsilon(3S) \gamma) = (2.13 \pm 0.04_{f_{-1.12}^{+0.75}} \pm 0.10_{\text{direct}} \pm 0.10_{h \rightarrow \gamma\gamma}) \cdot 10^{-9}$$

Branching Ratios in the SM

$$\text{BR}(h \rightarrow \rho^0 \gamma) = (1.68 \pm 0.02_f \pm 0.08_{h \rightarrow \gamma\gamma}) \cdot 10^{-5}$$

$$\text{BR}(h \rightarrow \omega \gamma) = (1.48 \pm 0.03_f \pm 0.07_{h \rightarrow \gamma\gamma}) \cdot 10^{-6}$$

$$\text{BR}(h \rightarrow \phi \gamma) = (2.31 \pm 0.03_f \pm 0.11_{h \rightarrow \gamma\gamma}) \cdot 10^{-6}$$

$$\text{BR}(h \rightarrow J/\psi \gamma) = (2.95 \pm 0.07_f \pm 0.06_{\text{direct}} \pm 0.14_{h \rightarrow \gamma\gamma}) \cdot 10^{-6}$$

$$\text{BR}(h \rightarrow \Upsilon(1S) \gamma) = (4.61 \pm 0.06_{f_{-1.21}^{+1.75}} \pm 0.22_{\text{direct}} \pm 0.22_{h \rightarrow \gamma\gamma}) \cdot 10^{-9}$$

$$\text{BR}(h \rightarrow \Upsilon(2S) \gamma) = (2.34 \pm 0.04_{f_{-0.99}^{+0.75}} \pm 0.11_{\text{direct}} \pm 0.11_{h \rightarrow \gamma\gamma}) \cdot 10^{-9}$$

$$\text{BR}(h \rightarrow \Upsilon(3S) \gamma) = (2.13 \pm 0.04_{f_{-1.12}^{+0.75}} \pm 0.10_{\text{direct}} \pm 0.10_{h \rightarrow \gamma\gamma}) \cdot 10^{-9}$$

The decays $h \rightarrow M\gamma$ are rare in the SM

Exclusive Radiative Decays $h \rightarrow MZ$

Alte, König and Neubert (2016)

Decay Topologies


Previous studies on some decay modes exist Isidori et al. (2014); Gao (2014);

Gonzalez-Alonso and Isidori (2014); Modak and Srivastava (2014)

Decay Topologies

Previous studies on some decay modes exist [Isidori et al. \(2014\)](#); [Gao \(2014\)](#);


[Gonzalez-Alonso and Isidori \(2014\)](#); [Modak and Srivastava \(2014\)](#)



Decay Topologies

Previous studies on some decay modes exist Isidori et al. (2014); Gao (2014);

Gonzalez-Alonso and Isidori (2014); Modak and Srivastava (2014)




The **diagram** $h \rightarrow Z(\gamma^* \rightarrow M)$ is **loop suppressed** but the off-shellness of the photon yields an **enhancement factor** $\left(\frac{m_h}{m_M}\right)^2$

Decay Topologies

Previous studies on some decay modes exist Isidori et al. (2014); Gao (2014);

Gonzalez-Alonso and Isidori (2014); Modak and Srivastava (2014)



The **diagram** $h \rightarrow Z(\gamma^* \rightarrow M)$ is **loop suppressed** but the off-shellness of the photon yields an **enhancement factor** $\left(\frac{m_h}{m_M}\right)^2$

Decay into **pseudoscalar (P)** or **vector mesons (V)**

Sensitivity to New-Physics Parameters

Pseudoscalar mesons (all units are MeV)

$$F^{\pi^0 Z} \approx 46.1 \kappa_Z \quad F^{\eta Z} \approx 27.7 \kappa_Z \quad F^{\eta' Z} \approx -33.8 \kappa_Z$$

Sensitivity to New-Physics Parameters

Pseudoscalar mesons (all units are MeV)

$$F^{\pi^0 Z} \approx 46.1 \kappa_Z \quad F^{\eta Z} \approx 27.7 \kappa_Z \quad F^{\eta' Z} \approx -33.8 \kappa_Z$$

κ_Z is already **strongly constrained**

Sensitivity to New-Physics Parameters

Pseudoscalar mesons (all units are MeV)

$$F^{\pi^0 Z} \approx 46.1 \kappa_Z \quad F^{\eta Z} \approx 27.7 \kappa_Z \quad F^{\eta' Z} \approx -33.8 \kappa_Z$$

κ_Z is already **strongly constrained**

Vector mesons

$$F_{\parallel}^{J/\psi Z} \approx 38.69 \kappa_Z - 1.75 + 0.73 \kappa_{\gamma Z}^{\text{eff}}$$

$$F_{\perp}^{J/\psi Z} \approx -294.8 + 123.1 \kappa_{\gamma Z}^{\text{eff}} + 38.69 \kappa_Z + 1.95 \kappa_c$$

The coefficient $\kappa_{\gamma Z}^{\text{eff}}$ parameterises **NP effects in the effective $h\gamma Z$ vertex** and **vanishes in the SM**

Branching Ratios as Functions of $\kappa_{\gamma Z}^{\text{eff}}$

The decay $h \rightarrow \gamma Z$ has not been observed and the **direct bounds on $\kappa_{\gamma Z}^{\text{eff}}$ and $\tilde{\kappa}_{\gamma Z}^{\text{eff}}$ are modest** CMS (2013)


$$\sqrt{|\kappa_{\gamma Z}^{\text{eff}} - 2.395|^2 + |\tilde{\kappa}_{\gamma Z}^{\text{eff}}|^2} < 7.2$$

Branching Ratios as Functions of $\kappa_{\gamma Z}^{\text{eff}}$

The decay $h \rightarrow \gamma Z$ has not been observed and the **direct bounds on $\kappa_{\gamma Z}^{\text{eff}}$ and $\tilde{\kappa}_{\gamma Z}^{\text{eff}}$ are modest** CMS (2013)

$$\sqrt{|\kappa_{\gamma Z}^{\text{eff}} - 2.395|^2 + |\tilde{\kappa}_{\gamma Z}^{\text{eff}}|^2} < 7.2$$

$\text{Br}(h \rightarrow \Upsilon(1S)Z)$ as a function of $\kappa_{\gamma Z}^{\text{eff}}$




Branching Ratios as Functions of $\kappa_{\gamma Z}^{\text{eff}}$

The decay $h \rightarrow \gamma Z$ has not been observed and the direct bounds on $\kappa_{\gamma Z}^{\text{eff}}$ and $\tilde{\kappa}_{\gamma Z}^{\text{eff}}$ are modest CMS (2013)

$$\sqrt{|\kappa_{\gamma Z}^{\text{eff}} - 2.395|^2 + |\tilde{\kappa}_{\gamma Z}^{\text{eff}}|^2} < 7.2$$

$\text{Br}(h \rightarrow \Upsilon(1S)Z)$ as a function of $\kappa_{\gamma Z}^{\text{eff}}$



$h \rightarrow \Upsilon(1S)Z$ can serve as complementary probes of $\kappa_{\gamma Z}^{\text{eff}}$

Branching Ratios in the Standard Model

Normalised to the **theory prediction for the Higgs width Γ_H** :

Decay mode	Branching ratio [10^{-6}]
$h \rightarrow \pi^0 Z$	$2.30 \pm 0.01_f \pm 0.09_{\Gamma_h}$
$h \rightarrow \eta Z$	$0.83 \pm 0.08_f \pm 0.03_{\Gamma_h}$
$h \rightarrow \eta' Z$	$1.24 \pm 0.12_f \pm 0.05_{\Gamma_h}$
$h \rightarrow \rho^0 Z$	$7.19 \pm 0.09_f \pm 0.28_{\Gamma_h}$
$h \rightarrow \omega Z$	$0.56 \pm 0.01_f \pm 0.02_{\Gamma_h}$
$h \rightarrow \phi Z$	$2.42 \pm 0.05_f \pm 0.09_{\Gamma_h}$
$h \rightarrow J/\psi Z$	$2.30 \pm 0.06_f \pm 0.09_{\Gamma_h}$
$h \rightarrow \Upsilon(1S) Z$	$15.38 \pm 0.21_f \pm 0.60_{\Gamma_h}$
$h \rightarrow \Upsilon(2S) Z$	$7.50 \pm 0.14_f \pm 0.29_{\Gamma_h}$
$h \rightarrow \Upsilon(3S) Z$	$5.63 \pm 0.10_f \pm 0.22_{\Gamma_h}$

Exclusive Radiative Decays $h \rightarrow MW$


Alte, König and Neubert (2016)

Decay Topologies

Idea: access **flavour-changing quark Yukawa** couplings in the decays $h \rightarrow MW$ Kagan et al. (2015)


Decay Topologies

Idea: access **flavour-changing quark Yukawa** couplings in the decays $h \rightarrow MW$ Kagan et al. (2015)



Decay Topologies

Idea: access **flavour-changing quark Yukawa** couplings in the decays $h \rightarrow MW$ Kagan et al. (2015)



For suitable decay modes, the **indirect contribution** can be **CKM suppressed**, while the **direct contribution** includes terms **enhanced by the top-quark mass**

Flavour-Changing Quark Yukawas in $h \rightarrow MW$

$$\text{Br}(h \rightarrow B^+ W^-) = 1.54 \cdot 10^{-10} \left(\kappa_W^2 + 427 \kappa_W \text{Re } Y_{ut} + 45615 |Y_{ut}|^2 \right)$$



$$\text{Br}(h \rightarrow B_c^+ W^-) = 8.21 \cdot 10^{-8} \left(\kappa_W^2 + 41 \kappa_W \text{Re } Y_{ct} + 413 |Y_{ct}|^2 \right)$$

Flavour-Changing Quark Yukawas in $h \rightarrow MW$

$$\text{Br}(h \rightarrow B^+ W^-) = 1.54 \cdot 10^{-10} (\kappa_W^2 + 427 \kappa_W \text{Re } Y_{ut} + 45615 |Y_{ut}|^2)$$

$$\text{Br}(h \rightarrow B_c^+ W^-) = 8.21 \cdot 10^{-8} (\kappa_W^2 + 41 \kappa_W \text{Re } Y_{ct} + 413 |Y_{ct}|^2)$$

Set $\kappa_W \rightarrow 1$ and $\text{Im}(Y_{qt}) \rightarrow 0$:





Flavour-Changing Quark Yukawas in $h \rightarrow MW$

$$\text{Br}(h \rightarrow B^+ W^-) = 1.54 \cdot 10^{-10} (\kappa_W^2 + 427 \kappa_W \text{Re } Y_{ut} + 45615 |Y_{ut}|^2)$$

$$\text{Br}(h \rightarrow B_c^+ W^-) = 8.21 \cdot 10^{-8} (\kappa_W^2 + 41 \kappa_W \text{Re } Y_{ct} + 413 |Y_{ct}|^2)$$

Set $\kappa_W \rightarrow 1$ and $\text{Im}(Y_{qt}) \rightarrow 0$:





Current **bounds from $\text{Br}(t \rightarrow qh)$** Buschmann et al. (2016): $\sqrt{|Y_{tq}|^2 + |Y_{qt}|^2} < 0.18$

Flavour-Changing Quark Yukawas in $h \rightarrow MW$

$$\text{Br}(h \rightarrow B^+ W^-) = 1.54 \cdot 10^{-10} (\kappa_W^2 + 427 \kappa_W \text{Re } Y_{ut} + 45615 |Y_{ut}|^2)$$

$$\text{Br}(h \rightarrow B_c^+ W^-) = 8.21 \cdot 10^{-8} (\kappa_W^2 + 41 \kappa_W \text{Re } Y_{ct} + 413 |Y_{ct}|^2)$$

Set $\kappa_W \rightarrow 1$ and $\text{Im}(Y_{qt}) \rightarrow 0$:



Current **bounds from $\text{Br}(t \rightarrow qh)$** Buschmann et al. (2016): $\sqrt{|Y_{tq}|^2 + |Y_{qt}|^2} < 0.18$

The dependence on Y_{qt} is strong

Conclusions

General feature of the decays $h \rightarrow MV$: rare in the SM

Conclusions

General feature of the decays $h \rightarrow MV$: **rare in the SM**

$h \rightarrow M\gamma$ can serve as **probes of light-quark Yukawa couplings**

Conclusions

General feature of the decays $h \rightarrow MV$: **rare in the SM**

$h \rightarrow M\gamma$ can serve as **probes of light-quark Yukawa couplings**

$h \rightarrow MZ$ can provide information about $\kappa_{\gamma Z}^{\text{eff}}$

Conclusions

General feature of the decays $h \rightarrow MV$: **rare in the SM**

$h \rightarrow M\gamma$ can serve as **probes of light-quark Yukawa couplings**

$h \rightarrow MZ$ can provide information about $\kappa_{\gamma Z}^{\text{eff}}$

$h \rightarrow MW$ exhibit a **strong sensitivity** to the **off-diagonal Yukawa couplings involving the top quark**